

Open Research Online

The Open University's repository of research publications and other research outputs

A review of volatiles in the Martian interior

Journal Item

How to cite:

Filiberto, Justin; Baratoux, David; Beaty, David; Breuer, Doris; Farcy, Benjamin J.; Grott, Matthias; Jones, John H.; Kiefer, Walter S.; Mane, Prakta; McCubbin, Francis M. and Schwenzer, Susanne P. (2016). A review of volatiles in the Martian interior. *Meteoritics and Planetary Science*, 51(11) 1935 -1958.

For guidance on citations see [FAQs](#).

© 2016 The Meteoritical Society



<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Version: Accepted Manuscript

Link(s) to article on publisher's website:
<http://dx.doi.org/doi:10.1111/maps.12680>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

A Review of Volatiles in the Martian Interior

Justin Filiberto^{1,2}, David Baratoux^{3,4}, David Beaty⁵, Doris Breuer⁶, Benjamin J. Farcy^{1,12},
Matthias Grott⁶, John H. Jones⁷, Walter S. Kiefer⁸, Prajkta Mane^{9,10}, Francis M. McCubbin¹¹,
and Susanne P. Schwenzer²

1 Department of Geology, Southern Illinois University, Carbondale, IL 62901, USA,

Filiberto@siu.edu

2 The Open University, Department of Environment, Earth & Ecosystems, Walton Hall, Milton
Keynes MK7 6AA, UK

3 Geosciences Environnement Toulouse, University of Toulouse, Institut de Recherche pour le
Développement & CNRS, 14 Avenue Edouard Belin 31 400, France

4 Institut Fondamental d'Afrique Noire Cheikh Anta Diop, Dakar, Senegal.

5 Mars Program Office, JPL/Caltech, Pasadena, CA 91109, USA

6 Institute of Planetary Research, German Aerospace Center (DLR), Berlin, Germany

7 XI-3, NASA Johnson Space Center Houston, TX 77058, USA

8 Lunar and Planetary Institute, Houston TX 77058, USA

9 School of Earth and Space Exploration, Arizona State University, Tempe, AZ 85287, USA

10 Center for Meteorite Studies, Arizona State University, Tempe, Arizona 85287-6004, USA

11 NASA Johnson Space Center, Mail Code XI2, 2101 NASA Parkway, Houston, TX 77058,
USA

12 Now at NASA Goddard Space Flight Center, 8800 Greenbelt Road, Greenbelt, MD 20771,
USA

Abstract

Multiple observations from missions to Mars have revealed compelling evidence for a volatile-rich Martian crust. A leading theory contends that eruption of basaltic magmas was the ultimate mechanism of transfer of volatiles from the mantle toward the surface after an initial outgassing related to the crystallization of a magma ocean. However, the concentrations of volatile species in ascending magmas and in their mantle source regions are highly uncertain. This work and this special issue of Meteoritics and Planetary Science summarize the key findings of the workshop on Volatiles in the Martian Interior (Nov. 3-4, 2014), the primary open questions related to volatiles in Martian magmas and their source regions, and the suggestions of the community at the workshop to address these open questions.

Introduction

The Workshop on Volatiles in the Martian Interior was held Nov. 3 – 4, 2014 at the Lunar and Planetary Institute in Houston, TX (<http://www.hou.usra.edu/meetings/volatiles2014/>) and gathered together scientists from diverse disciplines to discuss the state of knowledge of volatiles in the interior of Mars (see meeting report Filiberto et al. 2015). The goals of the workshop were to discuss the latest developments in the field, synthesize current knowledge, and identify the primary scientific questions that still need to be addressed. This review and the other papers published in this special issue of Meteoritics and Planetary Science will summarize the key findings of the workshop and primary open questions.

Three general processes determined the history, concentration, and present state of volatiles in the Martian interior: 1) accretion, 2) outgassing during magma ocean solidification, and 3) secondary degassing during later volcanism. Accretion of volatile-bearing material from the protoplanetary disk produced proto-Mars, which is thought to consist of building blocks that

formed rapidly (less than 5 million years) at $\sim 2 - 3$ AU and produced a planetary body that was more enriched in volatiles than Venus or Earth (e.g., Wänke 1981; Wänke and Dreibus 1988; Wänke 1991; Taylor 2001; Lunine et al. 2003; Dauphas and Pourmand 2011).

The halogen content of the Martian interior is thought to be ~ 2 times that of the Earth (Dreibus and Wänke 1985, 1987; Filiberto and Treiman 2009b; Taylor 2013). However, there is a large debate about the past and present water content of the Martian interior (e.g., McSween et al. 2001; Nekvasil et al. 2007; Filiberto and Treiman 2009b; McCubbin et al. 2010; McCubbin et al. 2012; Gross et al. 2013; Taylor 2013; Jones 2015). Further, models suggest that Mars potentially had higher water contents in the interior early in Mars' history (e.g., Médard and Grove 2006) and has subsequently lost volatiles from the interior during two later stages: early outgassing during differentiation (possibly during magma ocean solidification) and later degassing during volcanic eruptions. During crystallization and mantle melting, volatile elements would be enriched in the melt over the early crystallizing mineral phases (e.g., Aubaud et al. 2004; Dalou et al. 2012), depleting the mantle in volatile species. If there was no cap or early crust formation during the magma ocean period, the melt would rise to the surface by vigorous convection. At the surface of the magma ocean, the pressure could be low enough that saturation of some volatiles (specifically H_2O , CO_2 , Cl, and S) in the melt would be near zero (e.g., Dixon and Stolper 1995; Dixon et al. 1995; Webster et al. 1999; Gaillard and Scaillet 2014), causing volatile elements to outgas and potentially form an early atmosphere (e.g., Elkins-Tanton 2008; Hirschmann and Withers 2008). Secondary degassing during volcanic eruptions would further deplete the mantle of volatiles, as volatile elements are concentrated in magmas over residual mantle minerals. Without crustal recycling of elements back into the interior (e.g., Magna et al. 2015), the Martian mantle should become more depleted in volatile elements through time (e.g.,

McCubbin et al. 2008; Fraeman and Korenaga 2010; Grott et al. 2011; Morschhauser et al. 2011; Balta and McSween 2013; Jones 2015).

Approaches to quantifying and understanding the role of volatiles in planetary evolution

Efforts to determine concentrations of volatiles in the Martian interior have focused mainly on investigations of the Martian meteorites. Cosmochemical constraints on the volatile budget of Mars have been obtained from the bulk compositions of Martian meteorites. Mineral chemical constraints on the volatile budget of Martian magmas, and hence the Martian interior, have been obtained from studies of the hydrous phases apatite and amphibole in Martian meteorites. Isotopic studies of H, Cl, and S and noble gases have been used to investigate magmatic secondary volatile loss, variations within mantle source regions, and volatile evolution of the mantle, crust, and atmosphere. These studies have been combined with laboratory experiments and geophysical and geochemical models to constrain the volatile budget of Mars and to determine how the volatile content of the interior has evolved through time. Finally, the models produced have been constrained with planetary-scale observables including crustal thickness, surface and meteorite chemistry (specifically Cl, Th, and K), early formation of the crust, present-day volcanism, and no crustal recycling as implied by Martian meteorites (e.g., Magna et al. 2015). Here we discuss the state of the knowledge and open questions for different volatile species, place these in context of broader open questions about volatiles in the Martian interior, and make inferences about the data and/or samples that would be needed to address these open questions.

Current state of knowledge related to specific volatiles

Water, Fluorine, and Chlorine

Cosmochemical Constraints. The bulk water contents in shergottites are significantly lower than in comparable terrestrial basalts, suggesting that the present Martian interior is drier than the present terrestrial mantle (e.g., Carr and Wänke 1992; Watson et al. 1994a; Filiberto and Treiman 2009b; Jones 2015) and that degassing during the Hesperian and Amazonian may not have provided substantial water to the surface (e.g., Filiberto and Treiman 2009b; McCubbin et al. 2009; Filiberto et al. 2014c; Giesting and Filiberto 2016). However, the D/H ratio of the Martian mantle is similar to that of the terrestrial mantle (Owen et al. 1988; Watson et al. 1994b; Leshin et al. 1996; Leshin 2000; Boctor et al. 2003; Lunine et al. 2003; Usui et al. 2012, 2015; Mane et al. 2016), suggesting that sources of water on the two planets were similar. Models have suggested that the Martian mantle may have degassed its water early in its history (e.g., Médard and Grove 2006; Elkins-Tanton 2008; Grott et al. 2011) and lost its early water-rich atmosphere (e.g., Owen et al. 1988; Carr and Wänke 1992; Carr 1996; Lunine et al. 2003; Carr and Head 2010). However, there is a large debate about the pre-eruptive water content of Martian magmas and their source regions, with estimates as low as nearly-anhydrous (consistent with the bulk rock measurements) and as high as two weight percent dissolved water in the magma before eruption/emplacement (Treiman 1985; Mysen et al. 1998; Dann et al. 2001; McSween et al. 2001; Jones 2004; Herd et al. 2005; Nekvasil et al. 2007; Filiberto 2008; Nekvasil et al. 2009; McCubbin et al. 2010; McCubbin et al. 2012; Usui et al. 2012; Balta and McSween 2013; He et al. 2013; Giesting et al. 2015; Jones 2015; Usui et al. 2015). In comparing literature values for water abundances, it is important to recognize that some authors report results in terms of mantle source region abundance and others in terms of water dissolved in the magma. Because water is

incompatible, its abundance in the magma will typically be $\sim 10 - 20$ times larger than in the mantle source region, depending on the associated melt fraction. In order to constrain the water content of the Martian interior and investigate how the mantle may have changed through time, as well as investigating secondary processes like degassing and alteration, recent studies rely on hydrous minerals (apatite and amphibole) within Martian meteorites to determine the primary volatile contents of Martian magmas and their source regions (e.g., Treiman 1985; Watson et al. 1994b; Boctor et al. 2003; Greenwood et al. 2003; Patiño Douce and Roden 2006; McCubbin and Nekvasil 2008; Filiberto and Treiman 2009b; Patiño Douce et al. 2011; McCubbin et al. 2012; Gross et al. 2013; Filiberto et al. 2014c; Giesting et al. 2015; Howarth et al. 2015; McCubbin and Jones 2015; McCubbin et al. 2015).

Unlike water, since the mid 1980's we have had fairly good constraints on the halogen content, especially Cl, of Martian meteorites and hence their source regions (Dreibus and Wänke 1985, 1987). The Martian interior is thought to be $2 - 3$ times enriched in halogens compared with the terrestrial interior (Dreibus and Wänke 1985, 1987; Filiberto and Treiman 2009b; Taylor 2013; Filiberto et al. 2016). This is based on not just comparing the bulk chemistry of Martian meteorites and terrestrial basalts, but comparing ratios of volatile concentrations to a non-volatile element, thereby taking into account primary and secondary processing such as mantle melting, fractional crystallization, degassing, and alteration. The calculated Cl content in the shergottite source region is similar to the terrestrial enriched mantle, while the F content of the shergottite source region is similar to the terrestrial bulk mantle (Filiberto et al. 2016). In order to calculate the H_2O content of the Martian interior, Filiberto et al. (2016) relied on the H_2O/Cl ratio for Martian magmas from apatite and amphibole analyses (see below) and direct measurements of H_2O/Cl concentrations in glass melt inclusions from Usui et al. (2012; 2015).

Filiberto et al. (2016) calculated H₂O contents for the chassignite and nakhlite source region, as well as the source region for Yamato 980459 and LAR 06319. Filiberto et al. (2016) reported that the H₂O abundances of these sources are similar to the depleted terrestrial MORB source (Michael 1988; Saal et al. 2002). However, the large uncertainty in their calculations of an average source region for the Martian meteorites likely reflects heterogeneities in the Martian interior (such as the depleted, intermediate and enriched source regions, as well the chassignite-nakhlite source region) as explored by McCubbin et al. (2016) (see next section).

Volatile-bearing phases- Amphibole and Apatite. Instead of focusing on the bulk chemistry of the Martian meteorites, which may have been affected by secondary processes like degassing and alteration, many previous studies have focused on using the hydrous phases amphibole and apatite to constrain the Martian volatile history and budget (e.g., Treiman 1985; Watson et al. 1994b; Boctor et al. 2003; Greenwood et al. 2003; Patiño Douce and Roden 2006; McCubbin and Nekvasil 2008; Filiberto and Treiman 2009b; Patiño Douce et al. 2011; McCubbin et al. 2012; Gross et al. 2013; Giesting et al. 2015; Howarth et al. 2015; McCubbin et al. 2015; Howarth et al. 2016). The majority of apatites in Martian meteorites (**Figure 1**) are chlorine-rich (Patiño Douce and Roden 2006; Filiberto and Treiman 2009b; McCubbin et al. 2013; Howarth et al. 2015), which has previously led to the suggestion that Martian magmas are chlorine-rich with some magmas potentially having chlorine as the dominant volatile (or at least co-dominant with H₂O; e.g., Filiberto and Treiman, 2009b). However, apatite in Martian meteorites actually ranges from OH- bearing, F- apatite to almost pure end member Cl-apatite (**Figure 1**), which suggests a range in volatile contents of primary magmas with some being rich in H₂O (e.g., McCubbin and Nekvasil 2008; McCubbin et al. 2012; Gross et al. 2013; McCubbin et al. 2016). McCubbin et al. (2016) show a difference in the water content between the rare

earth element enriched and depleted sources (51 - 68 ppm vs 14 - 23 ppm H₂O; respectively) consistent with other observed geochemical differences. This work further supports the idea that water is heterogeneously distributed in the mantle. Further, by combining Martian apatite compositions with bulk S and Cl isotopic data, McCubbin et al. (2016) determined that most Martian meteorites have been affected by crustal contamination, and hence estimates of mantle volatile abundances from these samples is quite complicated.

Amphibole in Martian meteorites, when present, is typically high-Ti kaersutitic amphibole found in olivine-hosted melt inclusions (e.g., Floran et al. 1978; Treiman 1985). Kaersutitic amphibole has been analyzed for halogens and water in Zagami, Shergotty, Chassigny, and Northwest Africa 2737 (Treiman 1985; Watson et al. 1994a; Watson et al. 1994b; Beck et al. 2006; Treiman et al. 2007; McCubbin et al. 2010; He et al. 2013; Giesting et al. 2015). Kaersutite in Martian meteorites is typically oxy-amphibole and contains relatively low water (0.1 – 0.7 wt%) and Cl contents (~0.1 wt%) (Floran et al. 1978; Watson et al. 1994b; Treiman et al. 2007; McCubbin et al. 2010; Giesting et al. 2015). Calculated parental magmas for the kaersutite in the chassignites have H₂O/Cl ratios of 0.5 - 4.0 (Giesting et al. 2015), which are consistent with the parental magma volatile concentrations calculated from apatite analyses (H₂O/Cl ratios of 0.7 - 4.2) (McCubbin and Nekvasil 2008; Filiberto et al. 2016). This suggests a chlorine-rich, hydrous parental magma for the chassignites.

However, the nakhlites MIL 03346, 090030, 090032, 090136, and NWA 5790 do not contain kaersutite like the other Martian meteorites but instead contain potassic-chloro-hastingsite in melt inclusions (e.g., Sautter et al. 2006; McCubbin et al. 2009; McCubbin et al. 2013). Experimental evidence suggests that in order to crystallize such chloro-amphiboles, the activity of water in the melt inclusions must have been extremely low and the system must have

had adequate K and Fe to crystallize Cl-rich amphibole (Giesting and Filiberto 2016). This extreme Fe, K, and Cl enrichment may have been caused by exsolution of a saline [FeCl₂-rich (Bell and Simon 2011)] aqueous fluid from the late stage melt in the nakhlite pile that was then transported upward from a deeper source region in the nakhlite parent intrusion (McCubbin et al. 2013; Giesting and Filiberto 2016).

D/H ratio. The story of Martian D/H is somewhat in a state of flux. There are at least two water reservoirs on Mars, each having a distinct D/H ratio (**Figure 2**). The Martian mantle probably has a δD of $\leq -100\text{‰}$, whereas the atmospheric δD is $\sim +5000\text{‰}$ ($\delta D = [(D/H)_{\text{Sample}}/(D/H)_{\text{SMOW}}] - 1) \times 1000$; where SMOW is D/H ratio of Standard Mean Ocean Water = 0.0001559). Apart from these two, other reservoirs have been suggested (**Figure 2**). Here we will examine the two known reservoirs, the mantle and the atmosphere and briefly discuss a possible reservoir with a δD intermediate to the mantle and atmosphere end-members.

Particularly with respect to volatiles, the most promising sample of the Martian interior in our collections is Chassigny, as determined by the isotopic composition of Xe recorded in Chassigny bulk rock, which has a solar value (see section on Noble Gases) (Ott 1988). Therefore, it appears that the volatiles in Chassigny have not been in contact with the atmosphere, which has a different Xe-isotopic signature and fractionated D/H. Even so, the history of Martian mantle D/H, as mainly viewed from the Chassigny perspective, is still complex. Watson et al. (1994b) used the ion probe to investigate the D/H of olivine-hosted melt inclusions in Chassigny. One of these analyses yielded a δD of $\sim 4000\text{‰}$, which was in good agreement with Earth-based telescopic measurements of D/H in the Martian atmosphere (Owen et al. 1988). But Leshin et al. (1996) measured δD in stepwise thermal releases, up to 1000°C, and found that there was relatively little variation in δD between 200°C and 1000°C. δD of a

combusted Chassigny sample varied between -26‰ and -84‰. A Chassigny pyrolysis release gave δD values between -9 and -60‰. Clearly, these contrasting results from the same research group, using different analytical techniques, are not easily interpretable.

Another set of Martian meteorites that may come close to representing the Martian mantle signatures are the depleted, reduced shergottites, like Yamato 980459 and Tissint. Bockrath et al. (2003) carried out an ion probe study that reported D/H ratios in shergottites that were even lower than those of Leshin et al. (1996). However, in this case low δD values could possibly have been due to terrestrial contamination (e.g., epoxy in the thin section). Usui et al. (2012) measured δD in olivine-hosted melt-inclusion glasses from a primitive shergottite, Y980459. Usui et al. (2012) concluded, based on their results and those of Leshin et al. (1996), that the D/H of the Martian mantle was $\leq 275\text{‰}$ (**Figure 2**). This is consistent with the evidence that even Y980459 has experienced some crustal contamination (Peters et al. 2015). Chen et al. (2015) and Mane et al. (2016) measured impact melts and nominally anhydrous phases like merrillites and maskelynites in Tissint and concluded that the lowest observed δD value in Tissint ($\sim -100\text{‰}$) represents the Martian interior composition. The merrillites yielded the highest δD values (up to +3682 ‰) and is consistent with merrillite reacting with crustal H_2O on Mars to form a secondary whitlockite component, which can only occur on Mars at submagmatic temperatures (Hughes et al. 2008; McCubbin et al. 2014; Shearer et al. 2015). Although Tissint is not free of crustal contamination, these observations are consistent with the conclusions of Peters et al. (2015) that Tissint had experienced less crustal contamination than Y980459. Hydrogen isotopic composition of nominally anhydrous igneous mineral phases like olivines and pyroxenes in a nakhlite NWA 817 show the δD value -60 to -280 ‰ (Gillet et al. 2002). Similarly apatites from the Nakhla meteorite show a δD value of $\sim -100\text{‰}$ (Hallis et al. 2012). So, conservatively, we

recommend adopting a δD value of $\leq -100\%$ for the Martian mantle. In the sections on S and on Noble Gases, we will show that even Chassigny has experienced minor crustal contamination. Therefore, the lightest Chassigny δD value of Leshin et al. (1996) should probably be considered an upper limit.

It has long been known that the δD ratio of the Martian atmosphere is extremely high ($\sim 4000\%$; e.g., Owen et al., 1988); and these early measurements were later given support by other Earth-based telescopic/spectroscopic observations (e.g., Krasnopolsky and Feldman 2001). For comparison, the water of the Earth's oceans has a δD of 0% (by definition, e.g., Faure 1977), so the absolute D/H ratio of the Martian atmosphere is about 5 times higher than that of the Earth's oceans (**Figure 2**). Subsequently, δD in the Martian atmosphere has been measured directly by the Curiosity rover. The SAM instrument on Curiosity has a tunable laser spectrometer (TLS), which measured the δD of the Martian atmosphere to be $5000 \pm 1000\%$ (Webster et al. 2013). Therefore, we are confident that the D/H of the tenuous Martian atmosphere is 4000 - 6000% . The most likely reason for such a large D/H ratio, generally agreed upon (e.g., Pepin 1991) is preferential loss to space of H relative to D. Most recently, Villanueva et al. (2015) estimated that on Early Mars, water must have been abundant; from their measurements of atmospheric water and DHO across the globe, they calculated the ancient global equivalent layer was as high as 137 m. Mars has not had a dynamo-generated magnetic field since the end of the impact basin-forming epoch roughly 4 billion years ago (e.g., Lillis et al. 2008), and Jeans escape (i.e., loss of species that are not gravitationally bound) from the upper atmosphere and upper-atmospheric erosion by the solar wind can both contribute to preferential loss of lighter isotopes (e.g., Jakosky and Jones 1997).

Recently, Usui et al. (2015) and Mahaffy et al. (2015) have presented evidence for an

intermediate δD reservoir of 1000-2000‰ in Martian surficial materials. Usui et al. (2015) found a convergence of intermediate δD analyses from several Martian meteorites and Mahaffy et al. (2015) measured a similar range of δD for a Martian sediment. If these new analyses stand the test of time, it may be that we will have to consider multiple Martian δD reservoirs, not just the mantle and the atmosphere. Although at first glance, it may appear that mixing between mantle and surficial waters is the simplest solution for the origin of the intermediate reservoir, it is not altogether clear that this is true (Usui et al. 2015).

Consequently, Usui et al. (2015) presented two very different models for surficial Martian water. The first model assumed that there was a shallow, buried ice reservoir with an intermediate δD and that this reservoir dominated the surficial water budget. This reservoir was constrained to inefficiently communicate with the atmospheric reservoir. The atmospheric reservoir might also include water ice in the polar caps, which may sublime in response to upper atmospheric H loss. The second model of Usui et al. (2015) was that all surficial condensed waters have an intermediate δD and that the higher δD of the atmosphere has been maintained by a quasi-steady-state of sublimation and atmospheric loss. A third model, not considered by Usui et al. (2015) but explored by Greenwood et al. (2008), is that there was an early δD fractionation (by hydrodynamic escape?) followed by a slow, gradual increase in δD over geologic time. This model implicitly assumes that there is not enough surficial water exchanging with the atmosphere that the atmospheric δD is “buffered.” A probable consequence of this assumption is that the δD of all surficial reservoirs are changing with time.

Although there is uncertainty about how surficial δD reservoirs were established and how they are maintained, it is highly likely that some major δD fractionation occurred very early in Martian history. Of the various models described here, this early fractionation could have

produced a global surficial reservoir with a δD of 1000-5000‰.

Cl isotopes. Williams et al. (2016) and Sharp et al. (2016) investigate the Cl-isotopic composition of Martian meteorites to constrain Cl reservoirs in the Martian crust and interior and place the Cl isotopic composition of Mars in context with other planetary materials, the origin of early Solar System Cl-isotopic reservoirs, and the accretion of Mars.

Martian meteorites that are thought to have experienced crustal contamination have positive $\delta^{37}\text{Cl}$ values up to 8.6 ‰, which suggests a ^{37}Cl enriched crust (Williams et al. 2016; Sharp et al. 2016). Olivine-phyric shergottites and Shergotty have low $\delta^{37}\text{Cl}$ values (-4 to -2‰), which is thought to represent the primary Martian mantle value (-4 to -3‰). The low $\delta^{37}\text{Cl}$ values of the Martian mantle may represent the Cl-isotopic value of the primordial bulk composition of Mars inherited during accretion (Sharp et al. 2016). The other basaltic shergottites have $\delta^{37}\text{Cl}$ from 0 to +1 ‰ which may be due to mixing between a mantle source and a crustal source. Sharp et al. (2016) speculate that preferential loss of ^{35}Cl to space has resulted in a high $\delta^{37}\text{Cl}$ value for the Martian surface. However, recent results from MSL Curiosity show that the crustal chlorine isotope ratios at Gale Crater are more variable and negative in $\delta^{37}\text{Cl}$ than typically observed in SNC meteorite analyses, which was attributed to perchlorate fractionating chlorine isotopes (Farley et al. 2016).

Sulfur

Sulfur (in the form of sulfate) alteration and sedimentary minerals dominate what we know of the alteration mineralogy in the Martian crust, particularly in the Hesperian (e.g., Bibring et al. 2006; King and McLennan 2010). This has been used to suggest a change of alteration style from water-dominated alteration in the Noachian to S-dominated alteration in the

Hesperian, potentially caused by a massive outpouring of S-rich volcanism during the formation of Tharsis (e.g., Poulet et al. 2005; Bibring et al. 2006). Therefore, it is vital to understand the sulfur capacity of Martian magmas, as sulfur storage and transport between reservoirs, such as core, mantle, crust, and atmosphere, are tied to igneous processes (e.g., King and McSween 2005; Gaillard and Scaillet 2009; Richter et al. 2009; King and McLennan 2010; Ding et al. 2014; Gaillard and Scaillet 2014; Ding et al. 2015).

Martian meteorites contain up to ~ 3000 ppm S (King and McLennan 2010; Ding et al. 2015), compared with terrestrial basalts which typically contain up to ~ 2000 ppm S (e.g., Wallace and Edmonds 2011). Based on the sulfur content of Martian meteorites, the Martian interior contains 700 – 1000 ppm S (Ding et al. 2015) and 10.6 – 12.7 wt% S in the core (Stewart et al. 2007), although recent estimates based on Fe isotopes places an upper limit of approximately 8 wt.% S in the core (Shahar et al. 2015). The unknown S abundance, or other potential light elements in the core, has an important influence on the density and inferred radius of the Martian core (Mocquet et al. 2011); seismic measurements by the forthcoming Insight mission should improve our knowledge of the core radius and thus may indirectly help to constrain the abundance of S in the core.

Even though the Martian interior contains significant S, Martian magmas are not sulfide-saturated because they have relatively high S-solubility (up to ~ 4000 ppm S) (Richter et al. 2009; Ding et al. 2014), but may reach S-saturation after extensive crystallization (Baumgartner et al. 2015). If Martian magmas did not significantly outgassing sulfur, cumulates in the crust could be an important reservoir of magmatic sulfur (Ding et al. 2015). However, melting of the mantle and subsequent volcanism and degassing of sulfur-rich magmas could explain the abundance of surficial sulfate minerals (Gaillard and Scaillet 2009; Richter et al. 2009). Sulfur released as SO_2

into a volcanic gas from an H and C depleted melt would produce a strongly acidic brine (Gaillard and Scaillet 2009). But Martian sulfur processes are likely to be rather complex. For example, SO₂ degassing is often appealed to as a source for the ubiquitous S (sulfate and jarosite) in Martian soils (Poulet et al. 2005; Bibring et al. 2006; Gaillard and Scaillet 2009). However, the most primitive shergottites, as judged by their Mg# and ($\epsilon^{143}\text{Nd}$), have intrinsic oxygen fugacities that are near the iron-wustite buffer (e.g., Herd et al. 2002; Herd 2003). Under those redox conditions, almost all the S in a shergottite liquid will be speciated as FeS, not as sulfate, and iron sulfides are not particularly volatile — although any trace H₂S would be (Gaillard and Scaillet 2009; King and McLennan 2010; Métrich and Mandeville 2010). Therefore, before volcanism can degas significant amounts of oxidized S, the S must first be oxidized. Alternatively, sulfates could potentially form from weathering of sulfide-bearing basalts in a CO₂-rich environment which produced locally acidic environments (Dehouck et al. 2012). So at a minimum, the outgassing of volcanic S must be, at least, a two-stage process to oxidize the sulfur and form sulfates.

Sulfur Isotopes. Sulfur isotopes have recently been used to track crustal contamination in Martian meteorites and constrain whether the rare earth element enrichment seen in many Martian meteorites is due to crustal assimilation or different mantle source regions (Franz et al. 2014). Franz et al. (2014) showed that assimilation of sulfur-rich crustal material into Martian magmas was a common occurrence during the eruption and emplacement of the parental magmas to the Martian meteorites. Further, the range of sulfur-isotopic values in sulfides and sulfates in Martian meteorites cannot be explained with terrestrial analog processes, which would suggest a fundamental difference between sulfur chemical processes in the Martian atmosphere/crust and that of the Earth (Franz et al. 2014).

Carbon

The C content of the Martian interior is currently not well constrained with estimates varying over more than 3 orders of magnitude (e.g., Dasgupta et al. 2014). In addition, there is evidence that portions of the Martian mantle may be graphite saturated, which further complicates abundance estimates of C in the Martian mantle (e.g., Hirschmann and Withers 2008; Righter et al. 2008). Understanding the Martian carbon budget and any putative carbon cycle is vital because dissolved C will reduce the Martian mantle solidus, potentially drive volcanism, and is key for habitability of surficial environments (Steele et al. 2012a; Steele et al. 2012b; Dasgupta et al. 2014). Carbon solubility in Martian basaltic melts is small [0.34 – 2.12 wt% CO₂ for the Gusev Crater basalt “Humphrey” composition or 0.45 – 1.26 wt% CO₂ for a Yamato 908459 composition at ~1 – 3 GPa] and is relatively insensitive to bulk composition (Stanley et al. 2011; Stanley et al. 2012). However, carbon solubility in a magma changes as a function of temperature, pressure, oxygen fugacity, and H₂/H₂O fugacity (Hirschmann and Withers 2008; Ardia et al. 2013; Dasgupta et al. 2014; Li et al. 2014; Li et al. 2015). Carbon solubility in silicate melts generally increases with increasing temperature and oxygen fugacity; however, there is a complex interplay between $f\text{O}_2$ and $f\text{H}_2$ on the solubility of C in silicate melts. Carbon solubility typically increases with increasing $f\text{O}_2$ (e.g., Pawley et al. 1992; Holloway 1998; Stanley et al. 2014; Li et al. 2015); however, at extremely low $f\text{O}_2$ in H-rich systems, carbon solubility increases with increasing $f\text{H}_2$ and decreasing $f\text{O}_2$ (Hirschmann and Withers 2008; Ardia et al. 2013; Stanley et al. 2014; Li et al. 2015). The reason for this reversal is that the solution mechanism for C in silicate liquids changes as a function of H₂ fugacity, which in turn changes as a function of O₂ fugacity: as $f\text{O}_2$ decreases in the O-H system, the H₂/H₂O fugacity ratio increases (e.g., Elkins-Tanton and Grove 2011; Hirschmann et al. 2012;

Sharp et al. 2013). At high $f\text{O}_2$, carbon dissolves as carbonate in the melt, while at low $f\text{O}_2$ and high $f\text{H}_2$ carbon dissolves as C-H species (e.g., Mysen et al., 2009; Li et al., 2014; Ardia et al., 2013). Consequently, both the $f\text{O}_2$ and $f\text{H}_2$ of the Martian interior will control the primary C-species that would have been degassed from Martian magmas. Furthermore, it illustrates the potential link between low $f\text{O}_2$, high $f\text{H}_2$ magmas on Mars and the possible abiotic production of methane during episodes of Martian volcanism.

Carbon dissolution in a reduced magma ocean was likely limited (< 500 ppm C) and a magma-ocean post core formation was likely C-free with a core with ~ 1 wt.% C, if a single stage core-mantle equilibration model is applicable (Chi et al. 2014; Dasgupta et al. 2014). Assuming a magma ocean with low C contents and a reduced oxygen fugacity, degassing of a Martian magma ocean may have released methane to the atmosphere (Li et al. 2014); however degassed carbon alone may not be sufficient to maintain a warm surface of early Mars (e.g., Stanley et al. 2011; Li et al. 2014).

Reduced macromolecular carbon has been discovered in a number of Martian meteorites, both as inclusions within igneous mineral phases (Steele et al. 2012a; Agee et al. 2013), as well as macromolecular carbon and graphitic carbon associated with hydrothermally precipitated carbonate globules in Allan Hills 84001 (Steele et al. 2007; Steele et al. 2012b; Steele et al. 2016). Based on the petrologic context of this macromolecular carbon, it is interpreted as forming through abiotic synthesis under low $f\text{O}_2$, high $f\text{H}_2$ conditions. In fact, the presence of macromolecular carbon and polycyclic aromatic hydrocarbons in association with spinel inclusions in olivine from a number of Martian meteorites has been used as evidence to support that at least portions of the Martian mantle are graphite saturated (Steele et al. 2012a).

Noble Gases

Noble gases have been measured in the Martian atmosphere by the Viking lander (Owen et al. 1977) and Curiosity rover (Atreya et al. 2013; Mahaffy et al. 2013; Mahaffy et al. 2015; Conrad et al. 2016), in rock samples collected at Gale Crater by the Curiosity rover (Farley et al. 2014), and in the Martian meteorites (e.g., Swindle 2002 and references therein). In fact, noble gas measurements in shock melt glass pockets in EETA79001 provided the first clear evidence that the SNC meteorites are indeed from Mars (e.g., Bogard and Johnson 1983).

Noble gases in Martian meteorites represent a mixture of a variety of Martian and terrestrial components (**Figure 3**). On Mars, atmosphere, interior, and fractionated atmospheric components occur and specific isotopes are formed as radiogenic/nucleogenic products, including fission and spallation processes (e.g., Swindle 2002). During space travel, cosmogenic isotopes are formed. Once on Earth, terrestrial atmosphere, elementally fractionated in most cases, further complicate the picture (e.g., Mohapatra et al. 2009; Schwenzer et al. 2013).

From a noble gas perspective, Chassigny appears to be our best example of the interior (Ott 1988). It has a $^{129}\text{Xe}/^{132}\text{Xe}$ ratio of approximately 1, which means it contains no radiogenic ^{129}Xe and has a solar-like Xe-isotopic composition (Ott 1988). A second (^{136}Xe -enriched) reservoir has also been found in Chassigny (Mathew and Marti 2001). This ^{136}Xe -enriched component cannot be detected in bulk Chassigny analyses (e.g., Ott 1988), and can only be discerned by detailed, step-wise thermal releases (e.g., Mathew and Marti 2001). The second reservoir is thought to be an interior reservoir containing a contribution from Pu fission (e.g., Mathew and Marti 2001; Swindle 2002). This might be connected to the observation of two D/H reservoirs at different spatial resolution of the analyses (see above), and support the existence of inhomogeneity of the mantle volatile reservoir. Similarly, Ar isotopes are distinctive in the

Martian interior, but a second reservoir could exist (Schwenzer et al. 2007). Shergottite samples generally fall onto a trend between the atmospheric endmember measured by Viking (Owen et al. 1977), noble gases measured by Curiosity (Atreya et al. 2013; Conrad et al. 2016) , and a Martian interior component represented by Chassigny (e.g., Becker and Pepin 1984; Ott 1988).

Interestingly, at some point on Mars (possibly during crustal processes), elementally-fractionated atmospheric noble gas signatures were incorporated into the ALH 84001 and nakhlite Martian meteorites, complicating their noble gas signature and leading to a range of incorporation hypotheses (Ott 1988; Drake et al. 1994; Swindle et al. 1995; Gilmour et al. 1998; Mathew et al. 1998; Gilmour et al. 1999, 2001; Mathew and Marti 2001; Musselwhite and Swindle 2001; Mathew and Marti 2002; Swindle 2002; Mathew et al. 2003; Mathew and Marti 2005; Schwenzer and Ott 2006; Ott 2008; Mohapatra et al. 2009; Swindle et al. 2009; Cartwright et al. 2013). Further measurements of Martian rocks - preferably on Mars to avoid complications through later additions of isotopes, and careful investigations of fractionation effects, e.g., through experiments (Bullock et al. 2015; Schwenzer et al. 2016), are necessary to fully understand sources, pathways and sinks of the noble gases.

Effect on magma genesis and evolution of specific volatiles

Effect of water on magma genesis

Extensive work has investigated the effect of water on liquidus temperature, mantle solidus temperatures, and magma (basalt through granite) crystallization for terrestrial compositions (e.g., Burnham and Davis 1974; Mysen and Boettcher 1975; Danyushevsky 2001; Grove et al. 2002; Katz et al. 2003; Nekvasil et al. 2004; Feig et al. 2006; Grove et al. 2006;

Almeev et al. 2007; Médard and Grove 2008; Whitaker et al. 2008; Feig et al. 2010; Balta et al. 2011; Parman et al. 2011), with much less work focusing on the effect of water on magma crystallization and genesis for Martian compositions (e.g., Dann et al. 2001; Médard and Grove 2006; Nekvasil et al. 2007; Filiberto 2008; McCubbin et al. 2008; Nekvasil et al. 2009; Pommier et al. 2012). Water is known to depress basalt liquidus temperatures by breaking bridging oxygen bonds and thereby destabilizing plagioclase (e.g., Burnham and Davis 1974; Silver et al. 1990; Giordano et al. 2004; Mysen and Cody 2004). Water also has a large effect on the solidus temperature of the Martian mantle, which decreases from 975 °C at 1 GPa down to 800 °C at 3 GPa under water saturated conditions (e.g., Médard and Grove 2006). These solidus temperatures may be compared to the ones inferred for the dry primitive mantle of Mars, 1187 – 1220 °C at 1 GPa and 1396 – 1443 °C at 3 GPa (values estimated or calculated from Bertka and Holloway 1994; Filiberto and Dasgupta 2011; Matsukage et al. 2013; Kiefer et al., 2015; and Collinet et al. 2015). However, for concentrations below a few hundreds of ppm, water will have little effect on the position of the solidus despite having a potentially important effect on the rheology of the mantle.

Effect of water on magma crystallization and liquidus temperatures

In order to investigate the effect of H₂O on basaltic liquidus depression, experiments have been run on many different compositions at different pressures, temperatures, and using different techniques (e.g., Danyushevsky 2001; Katz et al. 2003; Feig et al. 2006; Almeev et al. 2007; Médard and Grove 2008; Feig et al. 2010). The resulting models of liquidus/solidus depression are all rather similar (Katz et al. 2003; Almeev et al. 2007; Médard and Grove 2008), except for that of Danyushevsky (2001) (**Figure 4**). Danyushevsky (2001) investigated the effect of small

amounts of H₂O on crystallization of terrestrial mid-ocean ridge and back-arc basin magmas and observed a greater liquidus depression (for the same water content) than the other studies, which suggests a bulk compositional dependence. Médard and Grove (2008) noted an effect of the bulk MgO content of the magma on the effect of H₂O on liquidus depression. They noted that magmas with higher MgO content have greater liquidus depression for the same weight percent water. Most Martian magmas have MgO contents similar to those of terrestrial magmas, which justifies using these models to explain Martian basalt genesis and crystallization. However, some Martian basalts have significantly higher MgO contents, and the parameterization of the effect of water on liquidus and solidus depression may break down for these compositions (**Figure 4**). In fact, preliminary results from hydrous experiments conducted on Martian meteorite Yamato 980459 with <0.5 wt% H₂O, 0.5 wt% H₂O, and 2 wt% H₂O (Norris and Herd 2006; Dalton et al. 2007; Draper 2007) have greater liquidus depression for the same water content than typical terrestrial basalts (**Figure 4**). It is important to note that the Yamato 980459 bulk composition has ~18 wt% MgO, significantly more than typical terrestrial basalts (< 12 wt%).

Effect of halogens on magma genesis

Experimental work on halogens in magmatic systems has mainly focused on its solubility in magmas, solution mechanisms, and effect on viscosity of evolved melt compositions for fluorine and basaltic systems for chlorine (e.g., Webster 1990; Webster 1997; Aiuppa et al. 2009; Filiberto and Treiman 2009a; Bell and Simon 2011; Dalou et al. 2012; Filiberto et al. 2014a; Dalou and Mysen 2015; Webster et al. 2015; Farcy et al. 2016). Halogens are known to have a large effect on the physical properties of a silicate melt; however, the effect of halogens on melt properties is largely dependent on melt chemistry (e.g., Metrich and Rutherford 1992; Dingwell

and Hess 1998; Zimova and Webb 2006; Baasner et al. 2013a; Filiberto et al. 2014a), which can result in wide-ranging implications for basalt genesis in both Earth and Mars. F and Cl are thought to form complexes with network modifying cations within a silicate melt, forming ion clusters that are distinct from the silicate network (Malinin et al. 1989; Webster and De Vivo 2002; Sandland et al. 2004; Evans et al. 2008; Filiberto and Treiman 2009a). F is thought to complex mainly with Mg in basaltic systems, but may also form complexes with Ca, K, and Al depending on the melt chemistry (Foley et al. 1986; Luth 1988b; Luth 1988a; Zeng and Stebbins 2000; Kiczinski et al. 2004; Filiberto et al. 2012). Studies of simplified silica-rich compositions (aluminosilicate glasses with and without sodium), have shown that fluorine preferentially complexes mainly with Al and Na but possibly Si as well (Schaller et al. 1992; Zeng et al. 1999; Zeng and Stebbins 2000; Liu and Nekvasil 2002; Liu and Tossell 2003; Mysen et al. 2004). Cl is thought to mainly form complexes with Fe^{2+} in basaltic systems, though it may complex with Ca or Mg as well (Metrich and Rutherford 1992; Webster and De Vivo 2002; Sandland et al. 2004; Zimova and Webb 2006; Filiberto and Treiman 2009a; Bell and Simon 2011; Filiberto et al. 2014a; Farcy et al. 2016).

These complexes affect silicate melts by decreasing their viscosity (though for certain bulk compositions, dissolved halogens may actually initially increase viscosity), by changing the polymerization of the silicate network (Dingwell et al. 1985; Dingwell 1989; Dingwell and Hess 1998; Giordano et al. 2004; Zimova and Webb 2006; Zimova and Webb 2007; Baasner et al. 2013a, b). As halogens form complexes with 2+ cations in the melt, the behavior of Fe and Mg partitioning between minerals and melt and therefore crystallizing mineral chemistry are affected (Filiberto and Treiman 2009a; Bell and Simon 2011; Filiberto et al. 2014a; Farcy et al. 2016). Finally, dissolved F and Cl will depress the liquidus temperature (Manning 1981; Luth 1988a;

Filiberto and Treiman 2009a; Filiberto et al. 2014a; Giehl et al. 2014; Farcy et al. 2016); however, the amount of liquidus depression is dependent on bulk composition (e.g., Filiberto et al. 2014a; Farcy et al. 2016).

Effect of moderately volatile elements on the mantle solidus

Although the primary focus of this review is on highly volatile elements and compounds, it is worth noting that moderately volatile elements (such as Na, K, and P) are also thought to be enriched in Mars relative to Earth (Dreibus and Wänke 1985; Taylor, 2013), and these enrichments have important effects on the Martian mantle solidus temperature. Sodium lowers the primitive mantle solidus by ~20 °C on Mars relative to Earth (Kiefer et al., 2015); its effect on the solidus will decline through Martian history as Na is partitioned from the mantle into the crust. Phosphorous and potassium will also act to lower the Martian solidus, although their effects will be smaller due to their smaller overall abundances in the Martian mantle. Kiefer et al. (2015) found that the elevated abundances of Na and K on Mars enhanced the time-integrated crustal production on Mars by 10-15% relative to models with Earth-like Na and K.

Geodynamical Modelling

The distribution of heat-producing elements (K, Th, U) and volatiles in crustal and mantle reservoirs is strongly connected with the thermal evolution of the planet. The concentration and distribution of K, Th, and U controls the heat budget and the vigor of convection, whereas water and other volatile elements control the rheology of the mantle, which also affects the vigor of convection. The efficiency of convection controls the heat transfer and heat loss, the thermal evolution of the mantle, and the rate of magmatic output or crustal

production, and therefore the formation of distinct reservoirs depleted or enriched in incompatible elements. Modeling of the thermal evolution of Mars addresses the coupling between these various processes with the objective of producing scenarios consistent with available observations, including petrological/geochemical/cosmochemical constraints, or constraints on the structure and dynamics of the Martian interior (e.g., crust thickness, existence of magnetic dynamos).

The thermal evolution of a terrestrial planet like Mars depends strongly on the rheology (viscosity) of the mantle material, which depends on temperature, pressure, and water content. Even a few tens of ppm water can be rheologically significant (Hirth and Kohlstedt 1996; Karato and Jung 1998; Mei and Kohlstedt 2000). Water, at an abundance of 200 ppm in the mantle source region, will reduce the viscosity by a factor of 36 relative to a nominally dry (10 ppm water) mantle (Mei and Kohlstedt 2000; Li et al. 2008), and the associated increase in convective vigor will increase the characteristic convective velocity by a factor of 11. As limiting cases, viscosities of 10^{19} and 10^{21} Pa s at a temperature of 1600 K and a pressure of 3 GPa are generally assumed to represent wet and dry mantle rheologies, respectively. The effects of water on the magma production rate are smaller. Comparison of magma production in dry mantle plume melting (Li and Kiefer 2007) and in water-undersaturated mantle plume melting (Kiefer and Li, this issue) shows that 200 ppm of water in the mantle source region increases the magma production rate by a factor of 3 under conditions appropriate to present-day Mars. Balta and McSween (2013) proposed that the shergottites represent melting from a hydrated mantle source, which would reduce the needed mantle melting temperature, while Filiberto and Dasgupta (2015) proposed that the shergottites represent localized dry mantle plume sources. Because water is incompatible in the minerals that dominate the mantle, it preferentially partitions into the

melt and escapes from the mantle to the crust. It is therefore crucial to include the effect of water on the solidus when modeling the coupled feedback loops between thermal evolution, mantle viscosity, magma production, and degassing.

Crustal production and implications for the heat budget and water concentration in the mantle

As a consequence of crustal formation, the mantle becomes depleted in water. Thermal evolution models typically predict a water depletion in the mantle by about 40-80 % (e.g., Fraeman and Korenaga 2010; Morschhauser et al. 2011; Sandu and Kiefer 2012). An example water degassing history is shown in **Figure 5**. The model (based on the “Mars Na+K” model in Figure 5 of Kiefer et al. (2015)) initially has 500 ppm of water in the primitive mantle. Two-thirds of this water degasses from the mantle, which is likely within the uncertainty of the geochemical constraints on volatile loss summarized below. Approximately 155 ppm of water remains in the mantle at present, consistent with the models of McCubbin et al. (2010; this issue) and Filiberto et al. (this issue) based on hydrous phases in Martian meteorites and Chen et al.’s (2015) study of hydrogen isotopes in the shergottite Tissint. Assuming that about 40 % of the water in the crust-forming melt is released into the atmosphere (Grott et al. 2011), a total of about 60-120 m of water is delivered to the surface for an initial amount of water in the mantle of 100 ppm. Most of the degassing occurs in the first billion years of Martian history, which is consistent with the concentration of valley networks in early Martian history (Hynek et al. 2010; Bouley et al. 2016), which might have contributed to an early Martian ocean (Clifford and Parker 2001; Perron et al. 2007). The crustal production for the model in Figure 5 (51 km globally averaged, Kiefer et al., 2015) is also consistent with observational constraints (Wieczorek and

Zuber, 2004). Some of the water that escapes from the mantle may remain trapped in the crust, resulting in a rheologically wet crust, which can lead to a mechanical decoupling of the crust and mantle (Grott and Breuer 2008, 2009). This may be necessary to explain the observed elastic lithosphere thickness in the Noachian with values smaller than about 20 km (e.g., McGovern et al. 2002; McGovern et al. 2004).

The water depletion in the mantle is consistent with the observation that about half of Mars' radioactive elements are in the crust (Norman 1999; McLennan 2001; Taylor et al. 2006; Hahn et al. 2011). Radioactive elements and H₂O are incompatible with similar partition coefficients ($D \sim 0.01$ -0.002) (Salters et al. 2002; Borg and Draper 2003; Aubaud et al. 2004), which implies that at least half of the total Mars water was released from the mantle – part of it is enriched in the crust and part of it degassed into the atmosphere. Note, however, that the concentration of radioactive heat sources in the crust is an estimate and in particular its vertical distribution is not well known. The results of the crustal radioactivity calculations were interpreted with the precaution that they would represent an upper limit if the Martian crust had developed a vertically heterogeneous chemistry with upward enrichment (Hahn et al. 2011) as in the case of the terrestrial continental crust.

However, the recent findings of feldspar-rich lithologies from orbital spectroscopy (Bandfield 2006; Carter and Poulet 2013; Wray et al. 2013), in-situ analyses at Gale crater (Sautter et al. 2014; Sautter et al. 2015), and within Martian meteorites (Filiberto et al. 2014b; Gross and Filiberto 2014; Santos et al. 2015) suggests that the structure of the Martian crust could be more complex than even a simple upward enrichment. The suggestion of silica-rich lithologies in ancient crustal material implies that lithologies enriched in heat-producing elements are not necessarily well exposed at the surface and may be eroded or buried by

Hesperian or Amazonian volcanic material, which is dominated by tholeiitic basalts (McSween et al. 2003; McSween et al. 2006). It has been indeed argued that a low-density component may be essentially buried (Pauer and Breuer 2008; Baratoux et al. 2014; Sautter et al. 2015). On the other hand, these supposed feldspar/silica-rich igneous rocks may represent plagioclase-enriched basaltic eruptive products and not silica-rich lithologies, and may represent localized outcrops that are not representative of a significant portion of the Martian crust (Rogers and Nekvasil 2015). Therefore, a better understanding of the architecture of the Martian crust, and of the relative contribution of light felsic intrusive components, or dense olivine-bearing tholeiitic basalts, is needed before K, Th, and U surface concentrations may be extrapolated to the entire crust and used to constrain the volatile and remaining heat budget of the crust and interior.

A much lower mantle water depletion (5-10 %) has been suggested by Hauck and Phillips (2002), which is likely because they used a partition coefficient for water (0.1) that is too large and thus underestimated the amount of water transport to the crust. A smaller depletion is also obtained when the change in mantle density due to crustal formation results in a layered mantle structure with an upper mantle being less dense and more depleted in incompatible elements than the lower mantle (Ruedas et al. 2013; Plesa and Breuer 2014). Subsequent melting of the depleted upper mantle results in a reduced outgassing rate in comparison to a well-mixed and homogeneous mantle and water is heterogeneously distributed. On the other hand, outgassing is strongly enhanced for efficient crustal recycling via delamination of the lower crust (Morschhauser et al. 2011; Ogawa and Yanagisawa 2012). Although crustal mixing is suggested to be inconsistent with the geochemical data (Mezger et al. 2013; Magna et al. 2015), this argument may not hold when Martian eclogite (e.g., Papike et al. 2013), i.e., dense rock resulting from high-pressure metamorphism of basalt, sinks to the core-mantle boundary to form a

separate reservoir. Nevertheless, this mechanism will not change the depletion of radioactive elements in the mantle but results in a more heterogeneous distribution of incompatible elements in the mantle with the co-existence of enriched and depleted reservoirs.

Heterogeneous mantle

Thermal evolution models consider a well-mixed homogeneous convecting mantle, while isotopic signatures in the Martian meteorites suggest the existence of distinct geochemical reservoirs that formed about 4.4-4.5 Ga ago and have not mixed since (e.g., Chen and Wasserburg 1986; Borg et al. 1997; Brandon et al. 2000; Halliday et al. 2001; Humayun et al. 2013; Bellucci et al. 2015; Nyquist et al. 2016). Distinct long-lived reservoirs are difficult to reconcile with a homogeneously convecting mantle, unless these reservoirs have been ‘captured’ early in the stagnant lid. Alternatively, fractional crystallization of a magma ocean has been suggested to explain the isotopic heterogeneities in the Martian meteorites (e.g., Elkins-Tanton et al. 2003; Elkins-Tanton 2008). In this scenario, the crystallization sequence of the magma ocean leads to a non-monotonic density increase and an exponential increase of incompatible elements towards the surface. Such a gravitationally unstable configuration implies a global mantle overturn resulting in a density stratification that is stable against thermal convection, assuming the most recent density profiles of fractional crystallization for Mars derived by Elkins-Tanton et al. (2005). It has been shown that such a scenario is not consistent with the subsequent volcanic history of Mars and also suggests extensive melting of the lower mantle (Plesa et al. 2014). A similar study, which instead assumed a much smaller density contrast was caused by fractional crystallization, suggests sufficient entrainment from the lower mantle after the overturn by thermal convection to permit the formation of small-scale domains of isotopically distinct

materials in the upper mantle (Scheinberg et al. 2014). However, a global mantle overturn would also suggest a more water-rich lower mantle and a depleted upper mantle after the overturn. Outgassing subsequent to the magma ocean crystallization is therefore likely to be very inefficient and a thin elastic lithosphere in the Noachian is difficult to obtain (see the review about the different thermal evolution scenarios of Mars; Breuer et al. 2016).

Degassing and atmospheric evolution

For a homogeneous mantle, most mantle water and CO₂ is released in the early evolution, i.e., during the first 500 Ma to 1 Ga when the bulk of the crust is also produced (**Figure 5**). A total of 0.9–1 bar CO₂ is outgassed during this time period if a mantle oxygen fugacity corresponding to one log₁₀ unit above the iron–wustite buffer is assumed (Hirschmann and Withers 2008; Grott et al. 2011). It is, however, controversial whether an atmosphere with a much higher pressure of several bars can result in surface temperatures reaching above the freezing point of water, necessary for the presence of long-standing global liquid surface water (Forget and Pierrehumbert 1997; Kasting 1997; Forget et al. 2013). In addition, atmospheric loss of H, due to thermal escape as a consequence of a high EUV flux of the young Sun during the first few hundred million years resulted in an entire loss of the early atmosphere (e.g., Tian et al. 2009; Lammer et al. 2013). Therefore, only volcanic activity and associated degassing after this initial loss phase may have contributed to the Martian atmosphere.

Constraints from the magnetic field

To explain the early magnetic field of Mars, a superheated core of a few hundred degrees has been commonly assumed to induce strong core cooling, sufficient to initiate a thermal

dynamo (e.g., Hauck and Phillips 2002; Breuer and Spohn 2003; Breuer and Spohn 2006; Morschhauser et al. 2011). This initial temperature distribution allows the generation of a thermal dynamo for up to a few hundred million years independent of the mantle rheology – although the duration of a magnetic field increases with the strength of mantle convection and thus the decrease in mantle viscosity. The dynamo then stops as a consequence of core cooling below a critical heat flow, i.e., the heat flow along the core adiabat (e.g., Breuer et al. 2010). The existence of a superheated core depends strongly on the core formation mechanism and is not well understood for Mars (Golabek et al. 2009). Sandu and Kiefer (2012) have shown that a thermal dynamo can be generated even without a superheated core if water is present because it strongly reduces the viscosity. The reduction in the convective vigor in the mantle due to mantle degassing reduces the heat flux out of the core and terminates a dynamo. Although many studies have suggested that the thermal effects of large impact basin formation are the cause of dynamo termination on Mars (e.g., Roberts et al. 2009), it is in fact possible to terminate the dynamo in the observed time period solely by mantle degassing and the associated reduction in heat flux out of the core. Moreover, it is possible that both internal and impact processes played a role in terminating the dynamo, because the impact mechanism works best if the heat flux out of the core has already been reduced to a level that is close to the critical flux for supporting dynamo generation.

Primary Open Questions

Discussion at the workshop focused on both our current state of knowledge (above) as well as the primary open questions about each volatile species and key higher level questions that are potentially answerable using either geophysics, additional studies of meteorite samples,

690 studies of samples returned by Mars sample return mission(s), or in situ measurements from
691 missions like the MSL Curiosity rover, Mars2020 rover, and/or ExoMars. As with the discussion
692 above about the current state of knowledge of volatiles in the Martian interior, here we discuss
693 the open questions for different volatile species before placing these in context of broader open
694 questions about volatiles in the Martian interior.

695 696 Water

697 Current studies have placed limits on the amount of water in the shergottite source
698 region, but there are still many critical unknowns about water in the Martian interior. Without
699 ancient samples of Martian basalts, it is difficult to constrain how water in the interior has
700 evolved through time. Further, Martian meteorites come from a limited number of unknown
701 locations on the Martian surface, so without context and with variable locations it is difficult to
702 constrain the heterogeneities of water in the Martian interior. This all leads to a critical unknown
703 of the input of magmatically degassed water to the Martian water cycle, which is vital to make a
704 habitable planet and explain the alteration minerals seen from orbit. Further, early
705 serpentinization of the Martian crust may help explain the remnant magnetic field intensity and
706 its distributions, and the present D/H ratio of the crust and atmosphere, but the input of water to
707 the surface and the global water concentrations need to be constrained. Therefore, what is the
708 role played by subsurface clathrates (or other crustal minerals) in trapping/releasing volatiles in
709 the Martian interior?

710 Further, how does water influence magma genesis, magma crystallization, and interior
711 dynamics? In particular, have volatiles aided in generating a diversity of Martian magmas, now
712 that new studies suggest possible occurrences of evolved rocks in the Martian crust? Does the

extensive work done on water in terrestrial systems apply to Martian magmas and the interior with significantly different chemistry? How does water (and other volatile species) influence Martian interior dynamics? Finally, how much water has been lost from the interior due to volcanism?

Halogens

We have a fairly good constraint on the concentration of chlorine in the Martian interior, but estimates for fluorine vary by an order of magnitude. We also have limited constraints on the halogen bearing phases in the Martian interior and how these halogen-bearing phases affect mantle properties including rheology, structure, and melting. Finally, what is the relative abundance of F:Cl:H₂O in Martian magmas and how do halogens and water combine to affect magma genesis, crystallization, and eruption?

Carbon

The largest unknown in the volatile budget of the Martian interior is: what is the bulk C concentration of Mars and is it higher or lower than the Earth? This is vital in order to establish the initial inventory of mantle carbon and understand the Martian carbon cycle from mantle to crust and finally atmosphere.

Isotopes and Noble Gases

The main question that came out of the workshop is the apparent non-correlation between H, S, and Cl isotopes. If all three are recording crustal contamination and secondary processes, then the analyses should correlate with each other, but the current data sets do not fully correlate.

Further, they do not correlate with the noble gas signatures, which also record secondary processes. How many reservoirs (e.g., more than one interior reservoir) do the isotopic and noble gas measurements represent? Which processes or factors (e.g., alteration mineral formation, T regime, redox, degassing) influence specific isotopic systems predominantly (but maybe not others to the same degree)? Finally, why do surface rocks not have the isotopic composition of the Martian atmosphere?

Geophysics

What is the relative importance of batch and fractional melting in Martian mantle plumes? Fractional melting tends to remove incompatible elements such as water early in the melting process, which means that water has less of an effect on the melt production rate in the fractional melting case. What role did mantle degassing and the associated reduction in mantle convection vigor play in terminating magnetic dynamo activity and producing discreet reservoirs in early Mars? How did/do volatiles influence the interior dynamics? Mantle water content has a large influence on the mantle solidus and mantle viscosity. Models of partial melting are quite dependent on interpretations of volatiles in the mantle. Will seismic observations from INSIGHT help resolve any of the issues discussed here?

Strategies moving forward

The community at the workshop concluded that in order to constrain bulk volatile and isotopic composition, and importantly the heterogeneity of the Martian interior, it is vital to increase the diversity of the sample collection, ideally including sophisticated in-situ measurements and through Mars sample return. Detailed experimental [petrologic studies](#) are

759 required to understand how volatiles affect mantle melting, magma genesis, and crustal
760 evolution. The results of these experiments should be taken into consideration for the
761 interpretation of global surface mineralogical or chemical trends in terms of planetary cooling
762 and/or mantle chemical evolution and heterogeneity. Finally, geophysical models will have to
763 include these experimental results along with the constraints from Martian samples.
764

Acknowledgements

The authors would like to thank the staff at the Lunar and Planetary Institute for providing logistical support for the workshop, the Mars Program Office for travel support for students BJF and PM, and the LPI for travel support for DB. The authors would like to thank T. Swindle and J. Mustard as well as the AE C.Goodrich for their constructive reviews which greatly improved the manuscript. The authors acknowledge support in part from NASA's Mars Fundamental Research Program grants NNX13AG35G to JF, NNX13AG44G to FMM, NNX14AJ94G to SPS and JF, and NNX14AG41G to WSK and NASA's Cosmochemistry Program grant #NNX11AK75G to M Wadhwa in support of PM. SPS also acknowledges support from an Open University Research Investment Fellowship.

776 REFERENCES

- 777 Agee, C.B., Wilson, N.V., McCubbin, F.M., Ziegler, K., Polyak, V.J., Sharp, Z.D., Asmerom,
778 Y., Nunn, M.H., Shaheen, R., Thiemens, M.H., 2013. Unique Meteorite from Early Amazonian
779 Mars: Water-Rich Basaltic Breccia Northwest Africa 7034. *Science* 339, 780-785.
- 780 Aiuppa, A., Baker, D.R., Webster, J.D., 2009. Halogens in volcanic systems. *Chemical Geology*
781 263, 1-18.
- 782 Almeev, R.R., Holtz, F., Koepke, J., Parat, F., Botcharnikov, R.E., 2007. The effect of H₂O on
783 olivine crystallization in MORB: Experimental calibration at 200 MPa. *American Mineralogist*
784 92, 670-674.
- 785 Ardia, P., Hirschmann, M.M., Withers, A.C., Stanley, B.D., 2013. Solubility of CH₄ in a
786 synthetic basaltic melt, with applications to atmosphere–magma ocean–core partitioning of
787 volatiles and to the evolution of the Martian atmosphere. *Geochimica et Cosmochimica Acta*
788 114, 52-71.
- 789 Atreya, S.K., Trainer, M.G., Franz, H.B., Wong, M.H., Manning, H.L.K., Malespin, C.A.,
790 Mahaffy, P.R., Conrad, P.G., Brunner, A.E., Leshin, L.A., Jones, J.H., Webster, C.R., Owen,
791 T.C., Pepin, R.O., Navarro-González, R., 2013. Primordial argon isotope fractionation in the
792 atmosphere of Mars measured by the SAM instrument on Curiosity and implications for
793 atmospheric loss. *Geophysical Research Letters* 40, 2013GL057763.
- 794 Aubaud, C., Hauri, E.H., Hirschmann, M.M., 2004. Hydrogen partition coefficients between
795 nominally anhydrous minerals and basaltic melts. *Geophys. Res. Lett.* 31, L20611.
- 796 Baasner, A., Schmidt, B.C., Webb, S.L., 2013a. Compositional dependence of the rheology of
797 halogen (F, Cl) bearing aluminosilicate melts. *Chemical Geology* 346, 172-183.
- 798 Baasner, A., Schmidt, B.C., Webb, S.L., 2013b. The effect of chlorine, fluorine and water on the
799 viscosity of aluminosilicate melts. *Chemical Geology* 357, 134-149.
- 800 Balta, J.B., Asimow, P.D., Mosenfelder, J.L., 2011. Hydrous, Low-carbon Melting of Garnet
801 Peridotite. *Journal of Petrology* 52, 2079-2105.
- 802 Balta, J.B., McSween, H.Y., 2013. Water and the composition of Martian magmas. *Geology* 41,
803 1115-1118.
- 804 Bandfield, J.L., 2006. Extended surface exposures of granitoid compositions in Syrtis Major,
805 Mars. *Geophysical Research Letters* 33.
- 806 Baratoux, D., Samuel, H., Michaut, C., Toplis, M.J., Monnereau, M., Wieczorek, M., Garcia, R.,
807 Kurita, K., 2014. Petrological constraints on the density of the Martian crust. *Journal of*
808 *Geophysical Research: Planets* 119, 1707-1727.

809 Baumgartner, R.J., Fiorentini, M.L., Baratoux, D., Micklethwaite, S., Sener, A.K., Lorand, J.P.,
810 McCuaig, T.C., 2015. Magmatic controls on the genesis of Ni–Cu±(PGE) sulphide
811 mineralisation on Mars. *Ore Geology Reviews* 65, Part 1, 400-412.

812 Beck, P., Barrat, J.A., Gillet, P., Wadhwa, M., Franchi, I.A., Greenwood, R.C., Bohn, M.,
813 Cotten, J., de Moortele, B.V., Reynard, B., 2006. Petrography and geochemistry of the
814 chassignite Northwest Africa 2737 (NWA 2737). *Geochimica et Cosmochimica Acta* 70, 2127-
815 2139.

816 Becker, R., Pepin, R., 1984. The case for a Martian origin of the shergottites: Nitrogen and noble
817 gases in EETA 79001. *Earth and Planetary Science Letters* 69, 225-242.

818 Bell, A.S., Simon, A., 2011. Experimental evidence for the alteration of the Fe³⁺/ΣFe of silicate
819 melt caused by the degassing of chlorine-bearing aqueous volatiles. *Geology* 39, 499-502.

820 Bellucci, J.J., Nemchin, A.A., Whitehouse, M.J., Humayun, M., Hewins, R., Zanda, B., 2015.
821 Pb-isotopic evidence for an early, enriched crust on Mars. *Earth and Planetary Science Letters*
822 410, 34-41.

823 Bertka, C.M., Holloway, J.R., 1994. Anhydrous Partial Melting of an Iron-Rich Mantle .1.
824 Subsolidus Phase Assemblages and Partial Melting Phase-Relations at 10 to 30 Kbar.
825 *Contributions to Mineralogy and Petrology* 115, 313-322.

826 Bibring, J.-P., Langevin, Y., Mustard, J.F., Poulet, F., Arvidson, R., Gendrin, A., Gondet, B.,
827 Mangold, N., Pinet, P., Forget, F., the, O.t., Berthe, M., Bibring, J.-P., Gendrin, A., Gomez, C.,
828 Gondet, B., Jouglet, D., Poulet, F., Soufflot, A., Vincendon, M., Combes, M., Drossart, P.,
829 Encrenaz, T., Fouchet, T., Mercurio, R., Belluci, G., Altieri, F., Formisano, V., Capaccioni, F.,
830 Cerroni, P., Coradini, A., Fonti, S., Korabiev, O., Kottsov, V., Ignatiev, N., Moroz, V., Titov, D.,
831 Zasova, L., Loiseau, D., Mangold, N., Pinet, P., Doute, S., Schmitt, B., Sotin, C., Hauber, E.,
832 Hoffmann, H., Jaumann, R., Keller, U., Arvidson, R., Mustard, J.F., Duxbury, T., Forget, F.,
833 Neukum, G., 2006. Global Mineralogical and Aqueous Mars History Derived from
834 OMEGA/Mars Express Data. *Science* 312, 400-404.

835 Boctor, N.Z., Alexander, C.M.O.D., Wang, J., Hauri, E., 2003. The sources of water in Martian
836 meteorites: clues from hydrogen isotopes. *Geochimica et Cosmochimica Acta* 67, 3971-3989.

837 Bogard, D.D., Johnson, P., 1983. Martian Gases in an Antarctic Meteorite. *Science* 221, 651-
838 654.

839 Borg, L.E., Draper, D.S., 2003. A petrogenetic model for the origin and compositional variation
840 of the martian basaltic meteorites. *Meteoritics & Planetary Science* 38, 1713-1731.

841 Borg, L.E., Nyquist, L.E., Taylor, L.A., Wiesmann, H., Shih, C.-Y., 1997. Constraints on
842 Martian differentiation processes from Rb · Sr and Sm · Nd isotopic analyses of the basaltic
843 shergottite QUE 94201. *Geochimica et Cosmochimica Acta* 61, 4915-4931.

844 Bouley, S., Baratoux, D., Matsuyama, I., Forget, F., Séjourné, A., Turbet, M., Costard, F., 2016.
845 Late Tharsis formation and implications for early Mars. *Nature* 531, 344-347.

846 Brandon, A., Walker, R., Morgan, J., Goles, G., 2000. Re-Os isotopic evidence for early
847 differentiation of the Martian mantle. *Geochimica et Cosmochimica Acta* 64, 4083-4095.

848 Breuer, D., Labrosse, S., Spohn, T., 2010. Thermal Evolution and Magnetic Field Generation
849 in Terrestrial Planets and Satellites. *Space Science Reviews* 152, 449-500.

850 Breuer, D., Spohn, T., 2003. Early plate tectonics versus single-plate tectonics on Mars:
851 Evidence from magnetic field history and crust evolution. *Journal of Geophysical Research:*
852 *Planets* 108, 10.1029/2002JE001999.

853 Breuer, D., Spohn, T., 2006. Viscosity of the Martian mantle and its initial temperature:
854 Constraints from crust formation history and the evolution of the magnetic field. *Planetary and*
855 *Space Science* 54, 153-169.

856 Brown, G.M., Peckett, A., 1977. Fluorapatites from the Skaergaard intrusion, East Greenland.
857 *Mineralogical Magazine* 41, 227-232.

858 Bullock, M.A., Schwenzer, S., Bridges, J., Chavez, C., Filiberto, J., Kelley, S., Miller, M.,
859 Moore, J., Smith, H., Swindle, T., Treiman, A., 2015. Noble gas fractionation during low
860 temperature alteration: an experimental approach. 46th Lunar and Planetary Science Conference,
861 Abstract #1235.

862 Burnham, C.W., Davis, F.A., 1974. The role of H₂O in silicate melts II: Thermodynamic and
863 phase relations in the system NaAlSi₃O₈-H₂O to 10 kilobars, 700° to 1000°C. *American Journal*
864 *of Science* 274, 902-940.

865 Carr, M.H., 1996. *Accretion and Evolution of Water, Water on Mars*. Oxford University Press,
866 New York, pp. 146-169.

867 Carr, M.H., Head, J.W., 2010. Geologic history of Mars. *Earth and Planetary Science Letters*
868 294, 185-203.

869 Carr, M.H., Wänke, H., 1992. Earth and Mars: Water inventories as clues to accretional histories.
870 *Icarus* 98, 61-71.

871 Carter, J., Poulet, F., 2013. Ancient plutonic processes on Mars inferred from the detection of
872 possible anorthositic terrains. *Nature Geosci* 6, 1008-1012.

873 Cartwright, J.A., Gilmour, J.D., Burgess, R., 2013. Martian fluid and Martian weathering
874 signatures identified in Nakhla, NWA 998 and MIL 03346 by halogen and noble gas analysis.
875 *Geochimica et Cosmochimica Acta* 105, 255-293.

876 Cartwright, J.A., Ott, U., Herrmann, S., Agee, C.B., 2014. Modern atmospheric signatures in 4.4
877 Ga Martian meteorite NWA 7034. *Earth and Planetary Science Letters* 400, 77-87.

878 Casey, J.F., Banerji, D., Zarian, P., 2007. Leg179 synthesis: geochemistry, stratigraphy, and
879 structure of gabbroic rocks drilled in ODP Hole 1105A, Southwest Indian Ridge., in: Casey, J.F.,

880 Miller, D.J. (Eds.), Proc. ODP, Sci. Results. Ocean Drilling Program, College Station, TX, pp. 1-
881 125. doi:110.2973/odp.proc.sr.2179.2001.2007.

882 Chen, J., Wasserburg, G., 1986. Formation ages and evolution of Shergotty and its parent planet
883 from U-Th-Pb systematics. *Geochimica et Cosmochimica Acta* 50, 955-968.

884 Chen, Y., Liu, Y., Guan, Y., Eiler, J.M., Ma, C., Rossman, G.R., Taylor, L.A., 2015. Evidence in
885 Tissint for recent subsurface water on Mars. *Earth and Planetary Science Letters* 425, 55-63.

886 Chi, H., Dasgupta, R., Duncan, M.S., Shimizu, N., 2014. Partitioning of carbon between Fe-rich
887 alloy melt and silicate melt in a magma ocean – Implications for the abundance and origin of
888 volatiles in Earth, Mars, and the Moon. *Geochimica et Cosmochimica Acta* 139, 447-471.

889 Clifford, S.M., Parker, T.J., 2001. The Evolution of the Martian Hydrosphere: Implications for
890 the Fate of a Primordial Ocean and the Current State of the Northern Plains. *Icarus* 154, 40-79.

891 Collinet, M., Médard, E., Charlier, B., Vander Auwera, J., Grove, T.L., 2015. Melting of the
892 primitive martian mantle at 0.5–2.2 GPa and the origin of basalts and alkaline rocks on Mars.
893 *Earth and Planetary Science Letters* 427, 83-94.

894 Conrad, P.G., Malespin, C.A., Franz, H.B., Pepin, R.O., trainer, M.G., Schwenzer, S.P., Atreya,
895 S.K., Freissinet, C., Jones, J.H., Manning, H., Owen, T., Pavlov, A.A., Wiens, R.C., Wong,
896 M.H., Mahaffy, P.R., 2016. In situ measurement of atmospheric krypton and xenon on Mars with
897 Mars Science Laboratory. *Earth and Planetary Science Letters*, in revision.

898 Coulson, I.M., Chambers, A.D., 1996. Patterns of zonation in rare-earth-bearing minerals in
899 nepheline syenites of the North Qoroq Center, South Greenland. *European Journal of Mineralogy*
900 34, 1163-1178.

901 Dalou, C., Koga, K., Shimizu, N., Boulon, J., Devidal, J.-L., 2012. Experimental determination
902 of F and Cl partitioning between lherzolite and basaltic melt. *Contributions to Mineralogy and*
903 *Petrology* 163, 591-609.

904 Dalou, C., Mysen, B.O., 2015. The effect of H₂O on F and Cl solubility and solution
905 mechanisms of in aluminosilicate melts at high pressure and high temperature. *American*
906 *Mineralogist* 100, 633-643.

907 Dalton, H.A., Sharp, T.G., Holloway, J.R., 2007. Investigation of the effects of water on a
908 Martian mantle composition. *Lunar and Planetary Science XXXVII*, Abstract # 2102.

909 Dann, J.C., Holzheid, A.H., Grove, T.L., McSween, H.Y., 2001. Phase equilibria of the
910 Shergotty meteorite: Constraints on pre-eruptive water contents of martian magmas and
911 fractional crystallization under hydrous conditions. *Meteoritics & Planetary Science* 36, 793-806.

912 Danyushevsky, L.V., 2001. The effect of small amounts of H₂O on crystallisation of mid-ocean
913 ridge and backarc basin magmas. *Journal of Volcanology and Geothermal Research* 110, 265-
914 280.

- 915 Dasgupta, R., Nelson, J.D., Chi, H., Ding, S., Li, Y., Duncan, M.S., Tsuno, K., 2014. Workshop
916 on Volatiles in the Martian Interior, Abstract #1012.
- 917 Dauphas, N., Pourmand, A., 2011. Hf-W-Th evidence for rapid growth of Mars and its status as
918 a planetary embryo. *Nature* 473, 489-492.
- 919 Dehouck, E., Chevrier, V., Gaudin, A., Mangold, N., Mathé, P.E., Rochette, P., 2012. Evaluating
920 the role of sulfide-weathering in the formation of sulfates or carbonates on Mars. *Geochimica et*
921 *Cosmochimica Acta* 90, 47-63.
- 922 Ding, S., Dasgupta, R., Lee, C.-T.A., Wadhwa, M., 2015. New bulk sulfur measurements of
923 Martian meteorites and modeling the fate of sulfur during melting and crystallization –
924 Implications for sulfur transfer from Martian mantle to crust–atmosphere system. *Earth and*
925 *Planetary Science Letters* 409, 157-167.
- 926 Ding, S., Dasgupta, R., Tsuno, K., 2014. Sulfur concentration of martian basalts at sulfide
927 saturation at high pressures and temperatures – Implications for deep sulfur cycle on Mars.
928 *Geochimica et Cosmochimica Acta* 131, 227-246.
- 929 Dingwell, D.B., 1989. Effect of fluorine on the viscosity of diopside liquid. *American*
930 *Mineralogist* 74, 333-338.
- 931 Dingwell, D.B., Hess, K.U., 1998. Melt viscosities in the system Na-Fe-Si-OF-Cl; contrasting
932 effects of F and Cl in alkaline melts. *American Mineralogist* 83, 1016-1021.
- 933 Dingwell, D.B., Scarfe, C.M., Cronin, D.J., 1985. The effect of fluorine on viscosities in the
934 system Na₂O-Al₂O₃-SiO₂; implications for phonolites, trachytes and rhyolites. *American*
935 *Mineralogist* 70, 80-87.
- 936 Dixon, J.E., Stolper, E.M., 1995. An Experimental Study of Water and Carbon Dioxide
937 Solubilities in Mid-Ocean Ridge Basaltic Liquids. Part II: Applications to Degassing *Journal of*
938 *Petrology* 36, 1633-1646.
- 939 Dixon, J.E., Stolper, E.M., Holloway, J.R., 1995. An Experimental Study of Water and Carbon
940 Dioxide Solubilities in Mid-Ocean Ridge Basaltic Liquids. *Journal of Petrology* 36, 1607-1631.
- 941 Drake, M.J., Swindle, T.D., Owen, T., Musselwhite, D.S., 1994. Fractionated martian
942 atmosphere in the nakhlites? *Meteoritics* 29, 854-859.
- 943 Draper, D.S., 2007. Water-undersaturated near-liquidus phase relations of Yamato 980459.
944 *Lunar and Planetary Science XXXVII*, Abstract #1447.
- 945 Dreibus, G., Wänke, H., 1985. Mars, a Volatile-Rich Planet. *Meteoritics* 20, 367-381.
- 946 Dreibus, G., Wänke, H., 1987. Volatiles on Earth and Mars - a Comparison. *Icarus* 71, 225-240.
- 947 Dymek, R.F., Owens, B.E., 2001. Petrogenesis of apatite-rich rocks (nelsonites and oxide-apatite
948 gabbro-norites) associated with massif anorthosites. *Economic Geology* 96, 797-815.

949 Elkins-Tanton, L.T., 2008. Linked magma ocean solidification and atmospheric growth for Earth
950 and Mars. *Earth and Planetary Science Letters* 271, 181-191.

951 Elkins-Tanton, L.T., Grove, T.L., 2011. Water (hydrogen) in the lunar mantle: Results from
952 petrology and magma ocean modeling. *Earth and Planetary Science Letters* 307, 173-179.

953 Elkins-Tanton, L.T., Hess, P.C., Parmentier, E.M., 2005. Possible formation of ancient crust on
954 Mars through magma ocean processes. *Journal of Geophysical Research* 110,
955 doi:10.1029/2005JE002480.

956 Elkins-Tanton, L.T., Parmentier, E.M., Hess, P.C., 2003. Magma ocean fractional crystallization
957 and cumulate overturn in terrestrial planets: Implications for Mars. *Meteoritics & Planetary
958 Science* 38, 1753-1771.

959 Evans, K.A., Mavrogenes, J.A., O'Neill, H.S., Keller, N.S., Jang, L.Y., 2008. A preliminary
960 investigation of chlorine XANES in silicate glasses. *Geochem. Geophys. Geosyst.* 9,
961 doi.10.1029/2008GC002157.

962 Farcy, B.J., Gross, J., Carpenter, P.K., Hicks, J., Filiberto, J., 2016. Effect of Cl on Near-
963 Liquidus Crystallization of Olivine-Phyric Shergottite NWA 6234 at 1 GPa: Implication for
964 volatile-induced melting of the Martian mantle. *Meteoritics and Planetary Science* this volume.

965 Farley, K.A., Malespin, C., Mahaffy, P., Grotzinger, J.P., Vasconcelos, P.M., Milliken, R.E.,
966 Malin, M., Edgett, K.S., Pavlov, A.A., Hurowitz, J.A., Grant, J.A., Miller, H.B., Arvidson, R.,
967 Beegle, L., Calef, F., Conrad, P.G., Dietrich, W.E., Eigenbrode, J., Gellert, R., Gupta, S.,
968 Hamilton, V., Hassler, D.M., Lewis, K.W., McLennan, S.M., Ming, D., Navarro-González, R.,
969 Schwenzer, S.P., Steele, A., Stolper, E.M., Sumner, D.Y., Vaniman, D., Vasavada, A., Williford,
970 K., Wimmer-Schweingruber, R.F., Team, t.M.S., 2014. In Situ Radiometric and Exposure Age
971 Dating of the Martian Surface. *Science* 343.

972 Farley, K.A., Martin, P., Archer Jr, P.D., Atreya, S.K., Conrad, P.G., Eigenbrode, J.L., Fairén,
973 A.G., Franz, H.B., Freissinet, C., Glavin, D.P., Mahaffy, P.R., Malespin, C., Ming, D.W.,
974 Navarro-Gonzalez, R., Sutter, B., 2016. Light and variable $^{37}\text{Cl}/^{35}\text{Cl}$ ratios in rocks from Gale
975 Crater, Mars: Possible signature of perchlorate. *Earth and Planetary Science Letters* 438, 14-24.

976 Faure, G., 1977. *Principles of isotope geology*.

977 Feig, S., Koepke, J., Snow, J., 2010. Effect of oxygen fugacity and water on phase equilibria of a
978 hydrous tholeiitic basalt. *Contributions to Mineralogy and Petrology* 160, 551-568.

979 Feig, S.T., Koepke, J., Snow, J.E., 2006. Effect of water on tholeiitic basalt phase equilibria: an
980 experimental study under oxidizing conditions. *Contributions to Mineralogy and Petrology* 152,
981 611-638.

982 Filiberto, J., 2008. Experimental Constraints on the Parental Liquid of the Chassigny Meteorite:
983 A Possible Link between the Chassigny Meteorite and a Gusev Basalt. *Geochimica et
984 Cosmochimica Acta* 72, 690-701.

985 Filiberto, J., Beaty, D., Kiefer, W.S., 2015. Volatiles in Mars: Constraints, questions, and future
986 directions. EOS, doi:10.1029/2015EO027375.

987 Filiberto, J., Dasgupta, R., 2011. Fe²⁺-Mg partitioning between olivine and basaltic melts:
988 Applications to genesis of olivine-phyric shergottites and conditions of melting in the Martian
989 interior. Earth and Planetary Science Letters 304, 527-537.

990 Filiberto, J., Dasgupta, R., 2015. Constraints on the depth and thermal vigor of melting in the
991 Martian mantle. Journal of Geophysical Research: Planets 120, 2014JE004745.

992 Filiberto, J., Dasgupta, R., Gross, J., Treiman, A., 2014a. Effect of chlorine on near-liquidus
993 phase equilibria of an Fe–Mg-rich tholeiitic basalt. Contributions to Mineralogy and Petrology
994 168, 1-13.

995 Filiberto, J., Gross, J., McCubbin, F.M., 2016. Constraints on the Water, Chlorine, and Fluorine
996 Content of the Martian Mantle. Meteoritics and Planetary Science, in press.

997 Filiberto, J., Gross, J., Trela, J., Ferré, E.C., 2014b. Gabbroic Shergottite Northwest Africa 6963:
998 An intrusive sample of Mars. American Mineralogist 99, 601-606.

999 Filiberto, J., Treiman, A.H., 2009a. The effect of chlorine on the liquidus of basalt: First results
1000 and implications for basalt genesis on Mars and Earth. Chemical Geology 263, 60-68.

1001 Filiberto, J., Treiman, A.H., 2009b. Martian magmas contained abundant chlorine, but little
1002 water. Geology 37, 1087-1090.

1003 Filiberto, J., Treiman, A.H., Giesting, P.A., Goodrich, C.A., Gross, J., 2014c. High-temperature
1004 chlorine-rich fluid in the martian crust: A precursor to habitability. Earth and Planetary Science
1005 Letters 401, 110-115.

1006 Filiberto, J., Wood, J., Dasgupta, R., Shimizu, N., Le, L., Treiman, A.H., 2012. Effect of fluorine
1007 on near-liquidus phase equilibria of an Fe–Mg rich basalt. Chemical Geology 312–313, 118-126.

1008 Floran, R.J., Prinz, M., Hlava, P.F., Keil, K., Nehru, C.E., Hinthorne, J.R., 1978. Chassigny
1009 Meteorite - Cumulate Dunite with Hydrous Amphibole-Bearing Melt Inclusions. Geochimica et
1010 Cosmochimica Acta 42, 1213-1229.

1011 Foley, S.F., Taylor, W.R., Green, D.H., 1986. The effect of fluorine on phase relationships in the
1012 system KAlSiO₄-Mg₂SiO₄-SiO₂ at 28 kbar and the solution mechanism of fluorine in silicate
1013 melts. Contributions to Mineralogy and Petrology 93, 46-55.

1014 Forget, F., Pierrehumbert, R.T., 1997. Warming early Mars with carbon dioxide clouds that
1015 scatter infrared radiation. Science 278, 1273-1276.

1016 Forget, F., Wordsworth, R., Millour, E., Madeleine, J.-B., Kerber, L., Leconte, J., Marcq, E.,
1017 Haberle, R.M., 2013. 3D modelling of the early martian climate under a denser CO₂
1018 atmosphere: Temperatures and CO₂ ice clouds. Icarus 222, 81-99.

- 1019 Fraeman, A.A., Korenaga, J., 2010. The influence of mantle melting on the evolution of Mars.
1020 *Icarus* 210, 43-57.
- 1021 Franz, H.B., Kim, S.-T., Farquhar, J., Day, J.M.D., Economos, R.C., McKeegan, K.D., Schmitt,
1022 A.K., Irving, A.J., Hoek, J., Iii, J.D., 2014. Isotopic links between atmospheric chemistry and the
1023 deep sulphur cycle on Mars. *Nature* 508, 364-368.
- 1024 Gaillard, F., Scaillet, B., 2009. The sulfur content of volcanic gases on Mars. *Earth and Planetary*
1025 *Science Letters* 279, 34-43.
- 1026 Gaillard, F., Scaillet, B., 2014. A theoretical framework for volcanic degassing chemistry in a
1027 comparative planetology perspective and implications for planetary atmospheres. *Earth and*
1028 *Planetary Science Letters* 403, 307-316.
- 1029 Giehl, C., Marks, M.W., Nowak, M., 2014. An experimental study on the influence of fluorine
1030 and chlorine on phase relations in peralkaline phonolitic melts. *Contributions to Mineralogy and*
1031 *Petrology* 167, 1-21.
- 1032 Giesting, P., Filiberto, J., 2016. The formation environment of potassic-chloro-hastingsite in the
1033 nakhlites MIL 03346 and pairs and NWA 5790: insights from terrestrial chloro-amphibole.
1034 *Meteoritics and Planetary Science*, in press.
- 1035 Giesting, P.A., Schwenzer, S.P., Filiberto, J., Starkey, N.A., Franchi, I.A., Treiman, A.H.,
1036 Tindle, A.G., Grady, M.M., 2015. Igneous and shock processes affecting chassignite amphibole
1037 evaluated using chlorine/water partitioning and hydrogen isotopes. *Meteoritics & Planetary*
1038 *Science* 50, 433-460.
- 1039 Gillet, P., Barrat, J., Deloule, E., Wadhwa, M., Jambon, A., Sautter, V., Devouard, B., Neuville,
1040 D., Benzerara, K., Lesourd, M., 2002. Aqueous alteration in the Northwest Africa 817 (NWA
1041 817) Martian meteorite. *Earth and Planetary Science Letters* 203, 431-444.
- 1042 Gilmour, J., Whitby, J., Turner, G., 1998. The Siting of Martian Xenon in Nakhla. *Meteoritics*
1043 *and Planetary Science Supplement* 33, 59.
- 1044 Gilmour, J., Whitby, J., Turner, G., 1999. Martian atmospheric xenon contents of Nakhla mineral
1045 separates: implications for the origin of elemental mass fractionation. *Earth and Planetary*
1046 *Science Letters* 166, 139-147.
- 1047 Gilmour, J., Whitby, J., Turner, G., 2001. Disentangling xenon components in Nakhla: Martian
1048 atmosphere, spallation and Martian interior. *Geochimica et Cosmochimica Acta* 65, 343-354.
- 1049 Giordano, D., Romano, C., Dingwell, D.B., Poe, B., Behrens, H., 2004. The combined effects of
1050 water and fluorine on the viscosity of silicic magmas. *Geochimica et Cosmochimica Acta* 68,
1051 5159-5168.
- 1052 Golabek, G., Gerya, T., Kaus, B., Ziethe, R., Tackley, P., 2009. Rheological controls on the
1053 terrestrial core formation mechanism. *Geochemistry, Geophysics, Geosystems* 10.

- 1054 Greenwood, J.P., Blake, R.E., Coath, C.D., 2003. Ion microprobe measurements of $^{18}\text{O}/^{16}\text{O}$
1055 ratios of phosphate minerals in the Martian meteorites ALH84001 and Los Angeles. *Geochimica*
1056 *et Cosmochimica Acta* 67, 2289-2298.
- 1057 Greenwood, J.P., Itoh, S., Sakamoto, N., Vicenzi, E.P., Yurimoto, H., 2008. Hydrogen isotope
1058 evidence for loss of water from Mars through time. *Geophys. Res. Lett.* 35.
- 1059 Gross, J., Filiberto, J., 2014. Granitic Compositions in Gabbroic Martian Meteorite NWA 6963
1060 and a Possible Connection to Felsic Compositions on the Martian Surface, Lunar and Planetary
1061 Institute Science Conference Abstracts, p. 1440.
- 1062 Gross, J., Filiberto, J., Bell, A.S., 2013. Water in the martian interior: Evidence for terrestrial
1063 MORB mantle-like volatile contents from hydroxyl-rich apatite in olivine-phyric shergottite
1064 NWA 6234. *Earth and Planetary Science Letters* 369–370, 120-128.
- 1065 Grott, M., Breuer, D., 2008. The evolution of the martian elastic lithosphere and implications for
1066 crustal and mantle rheology. *Icarus* 193, 503-515.
- 1067 Grott, M., Breuer, D., 2009. Implications of large elastic thicknesses for the composition and
1068 current thermal state of Mars. *Icarus* 201, 540-548.
- 1069 Grott, M., Morschhauser, A., Breuer, D., Hauber, E., 2011. Volcanic outgassing of CO_2 and
1070 H_2O on Mars. *Earth and Planetary Science Letters* 308, 391-400.
- 1071 Grove, T., Parman, S., Bowring, S., Price, R., Baker, M., 2002. The role of an H_2O -rich fluid
1072 component in the generation of primitive basaltic andesites and andesites from the Mt. Shasta
1073 region, N California. *Contributions to Mineralogy and Petrology* 142, 375-396.
- 1074 Grove, T.L., Chatterjee, N., Parman, S.W., Médard, E., 2006. The influence of H_2O on mantle
1075 wedge melting. *Earth and Planetary Science Letters* 249, 74-89.
- 1076 Hahn, B.C., McLennan, S.M., Klein, E.C., 2011. Martian surface heat production and crustal
1077 heat flow from Mars Odyssey Gamma-Ray spectrometry. *Geophys. Res. Lett.* 38, L14203.
- 1078 Halliday, A.N., Wänke, H., Birck, J.L., Clayton, R.N., 2001. The accretion, composition and
1079 early differentiation of Mars. *Space Science Reviews* 96, 197-230.
- 1080 Hallis, L.J., Taylor, G.J., Nagashima, K., Huss, G.R., 2012. Magmatic water in the martian
1081 meteorite Nakhla. *Earth and Planetary Science Letters* 359–360, 84-92.
- 1082 Hauck, S.A., Phillips, R.J., 2002. Thermal and crustal evolution of Mars. *Journal of Geophysical*
1083 *Research* 107, 5052.
- 1084 He, Q., Xiao, L., Hsu, W., Balta, J.B., McSween, H.Y., Liu, Y., 2013. The water content and
1085 parental magma of the second chassignite NWA 2737: Clues from trapped melt inclusions in
1086 olivine. *Meteoritics & Planetary Science* 48, 474-492.

1087 Herd, C.D.K., 2003. The oxygen fugacity of olivine-phyric martian basalts and the components
1088 within the mantle and crust of Mars *Meteoritics & Planetary Science* 38, 1711-1875.

1089 Herd, C.D.K., Borg, L.E., Jones, J.H., Papike, J.J., 2002. Oxygen fugacity and geochemical
1090 variations in the martian basalts: implications for martian basalt petrogenesis and the oxidation
1091 state of the upper mantle of Mars *Geochimica Et Cosmochimica Acta* 66, 2025-2036.

1092 Herd, C.D.K., Treiman, A.H., McKay, G.A., Shearer, C.K., 2005. Light lithophile elements in
1093 martian basalts: Evaluating the evidence for magmatic water degassing. *Geochimica et*
1094 *Cosmochimica Acta* 69, 2431-2440.

1095 Hirschmann, M.M., Withers, A.C., 2008. Ventilation of CO₂ from a reduced mantle and
1096 consequences for the early Martian greenhouse. *Earth and Planetary Science Letters* 270, 147-
1097 155.

1098 Hirschmann, M.M., Withers, A.C., Ardia, P., Foley, N.T., 2012. Solubility of molecular
1099 hydrogen in silicate melts and consequences for volatile evolution of terrestrial planets. *Earth*
1100 *and Planetary Science Letters* 345–348, 38-48.

1101 Hirth, G., Kohlstedt, D.L., 1996. Water in the oceanic upper mantle: implications for rheology,
1102 melt extraction and the evolution of the lithosphere. *Earth and Planetary Science Letters* 144, 93-
1103 108.

1104 Holloway, J.R., 1998. Graphite-melt equilibria during mantle melting: constraints on CO₂ in
1105 MORB magmas and the carbon content of the mantle. *Chemical Geology* 147, 89-97.

1106 Howarth, G., Pernet-Fisher, J., Bodnar, R., Taylor, L., 2015. Evidence for the exsolution of Cl-
1107 rich fluids in martian magmas: Apatite petrogenesis in the enriched lherzolithic shergottite
1108 Northwest Africa 7755. *Geochimica et Cosmochimica Acta*.

1109 Howarth, G.H., Liu, Y., Chen, Y., Pernet-Fisher, J.F., Taylor, L.A., 2016. Postcrystallization
1110 metasomatism in shergottites: Evidence from the paired meteorites LAR 06319 and LAR 12011.
1111 *Meteoritics & Planetary Science*, 10.1111/maps.12576.

1112 Hughes, J.M., Jolliff, B.L., Rakovan, J., 2008. The crystal chemistry of whitlockite and merrillite
1113 and the dehydrogenation of whitlockite to merrillite. *American Mineralogist* 93, 1300-1305.

1114 Humayun, M., Nemchin, A., Zanda, B., Hewins, R.H., Grange, M., Kennedy, A., Lorand, J.P.,
1115 Gopel, C., Fieni, C., Pont, S., Deldicque, D., 2013. Origin and age of the earliest Martian crust
1116 from meteorite NWA7533. *Nature* 503, 513-516.

1117 Hynek, B.M., Beach, M., Hoke, M.R.T., 2010. Updated global map of Martian valley networks
1118 and implications for climate and hydrologic processes. *Journal of Geophysical Research: Planets*
1119 115, n/a-n/a.

1120 Jakosky, B.M., Jones, J.H., 1997. The history of Martian volatiles. *Reviews of Geophysics* 35, 1-
1121 16.

- 1122 Jones, J.H., 2004. The Edge of Wetness: The Case for Dry Magmatism on Mars. Lunar and
1123 Planetary Science 35th, Abstract # 1798.
- 1124 Jones, J.H., 2015. Various aspects of the petrogenesis of the Martian shergottite meteorites.
1125 Meteoritics & Planetary Science 50, 674-690.
- 1126 Karato, S.-i., Jung, H., 1998. Water, partial melting and the origin of the seismic low velocity
1127 and high attenuation zone in the upper mantle. Earth and Planetary Science Letters 157, 193-207.
- 1128 Kasting, J.F., 1997. Warming early Earth and Mars. Science 276, 1213.
- 1129 Katz, R.F., Spiegelman, M., Langmuir, C.H., 2003. A new parameterization of hydrous mantle
1130 melting Geochemistry Geophysics Geosystems 4, 1073; DOI 10.1029/2002GC000433
- 1131 Kiczinski, T.J., Du, L.-S., Stebbins, J.F., 2004. F-19 NMR study of the ordering of high field
1132 strength cations at fluoride sites in silicate and aluminosilicate glasses. Journal of Non-
1133 Crystalline Solids 337, 142-149.
- 1134 Kiefer, W.S., Filiberto, J., Sandu, C., Li, Q., 2015. The effects of mantle composition on the
1135 peridotite solidus: Implications for the magmatic history of Mars. Geochimica et Cosmochimica
1136 Acta 162, 247-258.
- 1137 King, P.L., McLennan, S.M., 2010. Sulfur on Mars. Elements 6, 107-112.
- 1138 King, P.L., McSween, H.Y., 2005. Effects of H₂O, pH, and oxidation state on the stability of Fe
1139 minerals on Mars. Journal of Geophysical Research: Planets 110, n/a-n/a.
- 1140 Krasnopolsky, V.A., Feldman, P.D., 2001. Detection of molecular hydrogen in the atmosphere of
1141 Mars. Science 294, 1914-1917.
- 1142 Lammer, H., Chassefière, E., Karatekin, Ö., Morschhauser, A., Niles, P., Mousis, O., Odert, P.,
1143 Möstl, U., Breuer, D., Dehant, V., Grott, M., Gröller, H., Hauber, E., Pham, L., 2013. Outgassing
1144 History and Escape of the Martian Atmosphere and Water Inventory. Space Science Reviews
1145 174, 113-154.
- 1146 Leshin, L.A., 2000. Insights into martian water reservoirs from analyses of martian meteorite
1147 QUE94201. Geophysical Research Letters 27, 2017–2020.
- 1148 Leshin, L.A., Epstein, S., Stolper, E.M., 1996. Hydrogen isotope geochemistry of SNC
1149 meteorites. Geochimica et Cosmochimica Acta 60, 2635-2650.
- 1150 Li, Q., Kiefer, W.S., 2007. Mantle convection and magma production on present-day Mars:
1151 Effects of temperature-dependent rheology. Geophysical Research Letters 34,
1152 doi:10.1029/2007GL030544.
- 1153 Li, Y., Dasgupta, R., Tsuno, K., 2014. Workshop on Volatiles in the Martian Interior, Abstract
1154 #1016.

- 1155 Li, Y., Dasgupta, R., Tsuno, K., 2015. The effects of sulfur, silicon, water, and oxygen fugacity
1156 on carbon solubility and partitioning in Fe-rich alloy and silicate melt systems at 3 GPa and
1157 1600 °C: Implications for core–mantle differentiation and degassing of magma oceans and
1158 reduced planetary mantles. *Earth and Planetary Science Letters* 415, 54-66.
- 1159 Li, Z.-X.A., Lee, C.-T.A., Peslier, A.H., Lenardic, A., Mackwell, S.J., 2008. Water contents in
1160 mantle xenoliths from the Colorado Plateau and vicinity: Implications for the mantle rheology
1161 and hydration-induced thinning of continental lithosphere. *Journal of Geophysical Research:*
1162 *Solid Earth* 113, n/a-n/a.
- 1163 Lillis, R.J., Frey, H.V., Manga, M., 2008. Rapid decrease in Martian crustal magnetization in the
1164 Noachian era: Implications for the dynamo and climate of early Mars. *Geophysical Research*
1165 *Letters* 35, n/a-n/a.
- 1166 Liu, Y., Comodi, P., 1993. Some aspects of the crystal-chemistry of apatites. *Mineralogical*
1167 *Magazine* 57, 709-719.
- 1168 Liu, Y., Nekvasil, H., 2002. Si-F bonding in aluminosilicate glasses: Inferences from ab initio
1169 NMR calculations. *American Mineralogist* 87, 339-346.
- 1170 Liu, Y., Tossell, J., 2003. Possible Al–F Bonding Environment in Fluorine-Bearing Sodium
1171 Aluminosilicate Glasses: From Calculation of 19F NMR Shifts. *The Journal of Physical*
1172 *Chemistry B* 107, 11280-11289.
- 1173 Lunine, J.I., Chambers, J., Morbidelli, A., Leshin, L.A., 2003. The origin of water on Mars.
1174 *Icarus* 165, 1-8.
- 1175 Luth, R.W., 1988a. Effects of F on phase equilibria and liquid structure in the system NaAlSiO₄-
1176 CaMgSi₂O₆-SiO₂. *American Mineralogist* 73, 306-312.
- 1177 Luth, R.W., 1988b. Raman spectroscopic study of the solubility mechanisms of F in glasses in
1178 the system CaO-CaF-SiO. *American Mineralogist* 73, 297-305.
- 1179 Magna, T., Gussone, N., Mezger, K., 2015. The calcium isotope systematics of Mars. *Earth and*
1180 *Planetary Science Letters* 430, 86-94.
- 1181 Mahaffy, P.R., Conrad, P.G., Team, M.S., 2015. Volatile and Isotopic Imprints of Ancient Mars.
1182 *Elements* 11, 51-56.
- 1183 Mahaffy, P.R., Webster, C.R., Atreya, S.K., Franz, H., Wong, M., Conrad, P.G., Harpold, D.,
1184 Jones, J.J., Leshin, L.A., Manning, H., Owen, T., Pepin, R.O., Squyres, S., Trainer, M., Team,
1185 M.S., 2013. Abundance and Isotopic Composition of Gases in the Martian Atmosphere from the
1186 Curiosity Rover. *Science* 341, 263-266.
- 1187 Malinin, S.D., Kravchuk, I.F., Delbove, F., 1989. Chloride distribution between phases in
1188 hydrated and dry chloride-aluminosilicate melt systems as a function of phase composition.
1189 *Geochemistry International* 26, 32-38.

1190 Mane, P., Hervig, H., Wadhwa, M., Garvie, L.A.J., Balta, J.B., McSween, H.Y.J., 2016.
 1191 Hydrogen isotopic composition of the martian mantle inferred from the newest martian meteorite
 1192 fall Tissint. *Meteoritics and Planetary Science* this volume.

1193 Manning, D.A.C., 1981. The effect of fluorine on liquidus phase relationships in the system Qz-
 1194 Ab-Or with excess water at 1 kb. *Contributions to Mineralogy and Petrology* 76, 206-215.

1195 Mathew, K., Kim, J., Marti, K., 1998. Martian atmospheric and indigenous components of xenon
 1196 and nitrogen in SNC meteorites. *Meteoritics and Planetary Science* 33, 655-664.

1197 Mathew, K., Marti, K., 2002. Martian atmospheric and interior volatiles in the meteorite Nakhla.
 1198 *Earth and Planetary Science Letters* 199, 7-20.

1199 Mathew, K., Marti, K., 2005. Evolutionary trends in volatiles of the nakhlite source region of
 1200 Mars. *Journal of Geophysical Research: Planets* (1991–2012) 110.

1201 Mathew, K., Marty, B., Marti, K., Zimmermann, L., 2003. Volatiles (nitrogen, noble gases) in
 1202 recently discovered SNC meteorites, extinct radioactivities and evolution. *Earth and Planetary*
 1203 *Science Letters* 214, 27-42.

1204 Mathew, K.J., Marti, K., 2001. Early evolution of Martian volatiles: Nitrogen and noble gas
 1205 components in ALH84001 and Chassigny. *Journal of Geophysical Research: Planets* 106, 1401-
 1206 1422.

1207 Matsukage, K.N., Nagayo, Y., Whitaker, M.L., Takahashi, E., Toshisuke, K., 2013. Melting of
 1208 the Martian mantle from 1.0 to 4.5GPa. *Journal of Mineralogical and Petrological Sciences* 108,
 1209 201-214.

1210 McCubbin, F.M., Boyce, J.W., Srinivasan, P., Elardo, S.M., Santos, A.R., Filiberto, J., Shearer
 1211 Jr, C.K., 2016. Heterogeneous distribution of H₂O in the martian interior: Implications for the
 1212 abundances of H₂O in depleted and enriched mantle sources. *Meteoritics and Planetary Science*,
 1213 in press.

1214 McCubbin, F.M., Elardo, S.M., Shearer, C.K., Smirnov, A., Hauri, E.H., Draper, D.S., 2013. A
 1215 petrogenetic model for the comagmatic origin of chassignites and nakhlites: Inferences from
 1216 chlorine-rich minerals, petrology, and geochemistry. *Meteoritics & Planetary Science* 48, 819-
 1217 853.

1218 McCubbin, F.M., Hauri, E.H., Elardo, S.M., Vander Kaaden, K.E., Wang, J., Shearer, C.K.,
 1219 2012. Hydrous melting of the martian mantle produced both depleted and enriched shergottites.
 1220 *Geology* 40, 683-686.

1221 McCubbin, F.M., Jones, R.H., 2015. Extraterrestrial Apatite: Planetary Geochemistry to
 1222 Astrobiology. *Elements* 11, 183-188.

1223 McCubbin, F.M., Nekvasil, H., 2008. Maskelynite-hosted apatite in the Chassigny meteorite:
 1224 Insights into late-stage magmatic volatile evolution in martian magmas. *American Mineralogist*
 1225 93, 676-684.

- 1226 McCubbin, F.M., Nekvasil, H., Harrington, A.D., Elardo, S.M., Lindsley, D.H., 2008.
 1227 Compositional diversity and stratification of the Martian crust: Inferences from crystallization
 1228 experiments on the picrobasalt Humphrey from Gusev Crater, Mars,. *Journal of Geophysical*
 1229 *Research* 113, doi:10.1029/2008JE003165.
- 1230 McCubbin, F.M., Shearer, C.K., Burger, P.V., Hauri, E.H., Wang, J., Elardo, S.M., Papike, J.J.,
 1231 2014. Volatile abundances of coexisting merrillite and apatite in the martian meteorite Shergotty:
 1232 Implications for merrillite in hydrous magmas. *American Mineralogist* 99, 1347-1354.
- 1233 McCubbin, F.M., Smirnov, A., Nekvasil, H., Wang, J., Hauri, E., Lindsley, D.H., 2010. Hydrous
 1234 magmatism on Mars: A source of water for the surface and subsurface during the Amazonian.
 1235 *Earth and Planetary Science Letters* 292, 132-138.
- 1236 McCubbin, F.M., Tosca, N.J., Smirnov, A., Nekvasil, H., Steele, A., Fries, M., Lindsley, D.H.,
 1237 2009. Hydrothermal jarosite and hematite in a pyroxene-hosted melt inclusion in martian
 1238 meteorite Miller Range (MIL) 03346: Implications for magmatic-hydrothermal fluids on Mars.
 1239 *Geochimica et Cosmochimica Acta* 73, 4907-4917.
- 1240 McCubbin, F.M., Vander Kaaden, K.E., Tartèse, R., Boyce, J.W., Mikhail, S., Whitson, E.S.,
 1241 Bell, A.S., Anand, M., Franchi, I.A., Wang, J., Hauri, E.H., 2015. Experimental investigation of
 1242 F, Cl, and OH partitioning between apatite and Fe-rich basaltic melt at 1.0-1.2 GPa and 950-
 1243 1000 °C. *American Mineralogist* 100, 1790-1802.
- 1244 McGovern, P., Smith, J., Morgan, J., Bulmer, M., 2004. Olympus Mons aureole deposits: New
 1245 evidence for a flank failure origin. *Journal of Geophysical Research: Planets* (1991–2012) 109.
- 1246 McGovern, P.J., Solomon, S.C., Smith, D.E., Zuber, M.T., Simons, M., Wieczorek, M.A.,
 1247 Phillips, R.J., Neumann, G.A., Aharonson, O., Head, J.W., 2002. Localized gravity/topography
 1248 admittance and correlation spectra on Mars: Implications for regional and global evolution.
 1249 *Journal of Geophysical Research: Planets* 107, 19-11-19-25.
- 1250 McLennan, S.M., 2001. Crustal heat production and the thermal evolution of Mars. *Geophysical*
 1251 *Research Letters* 28, 4019-4022.
- 1252 McSween, H.Y., Grove, T.L., Lentz, R.C., Dann, J.C., Holzheid, A.H., Riciputi, L.R., Ryan,
 1253 J.G., 2001. Geochemical evidence for magmatic water within Mars from pyroxenes in the
 1254 Shergotty meteorite. *Nature* 409, 487-490.
- 1255 McSween, H.Y., Grove, T.L., Wyatt, M.B., 2003. Constraints on the composition and
 1256 petrogenesis of the Martian crust. *Journal of Geophysical Research-Planets* 108.
- 1257 McSween, H.Y., Wyatt, M.B., Gellert, R., Bell, J.F., Morris, R.V., Herkenhoff, K.E., Crumpler,
 1258 L.S., Milam, K.A., Stockstill, K.R., Tornabene, L.L., Arvidson, R.E., Bartlett, P., Blaney, D.,
 1259 Cabrol, N.A., Christensen, P.R., Clark, B.C., Crisp, J.A., Des Marais, D.J., Economou, T.,
 1260 Farmer, J.D., Farrand, W., Ghosh, A., Golombek, M., Gorevan, S., Greeley, R., Hamilton, V.E.,
 1261 Johnson, J.R., Joliff, B.L., Klingelhöfer, G., Knudson, A.T., McLennan, S., Ming, D., Moersch,
 1262 J.E., Rieder, R., Ruff, S.W., Schröder, C., de Souza, P.A., Squyres, S.W., Wänke, H., Wang, A.,
 1263 Yen, A., Zipfel, J., 2006. Characterization and petrologic interpretation of olivine-rich basalts at

- 1264 Gusev Crater, Mars. *Journal of Geophysical Research-Planets* 111, E02510,
1265 doi:1029/2005E02477.
- 1266 Médard, E., Grove, T., 2008. The effect of H₂O on the olivine liquidus of basaltic melts:
1267 experiments and thermodynamic models. *Contributions to Mineralogy and Petrology* 155, 417-
1268 432.
- 1269 Médard, E., Grove, T.L., 2006. Early hydrous melting and degassing of the Martian interior.
1270 *Journal of Geophysical Research* 111, E11003.
- 1271 Mei, S., Kohlstedt, D., 2000. Influence of water on plastic deformation of olivine aggregates: 2.
1272 Dislocation creep regime. *Journal of Geophysical Research: Solid Earth* (1978–2012) 105,
1273 21471-21481.
- 1274 Métrich, N., Mandeville, C.W., 2010. Sulfur in Magmas. *Elements* 6, 81-86.
- 1275 Métrich, N., Rutherford, M.J., 1992. Experimental study of chlorine behavior in hydrous silicic
1276 melts. *Geochimica et Cosmochimica Acta* 56, 607-616.
- 1277 Mezger, K., Debaille, V., Kleine, T., 2013. Core formation and mantle differentiation on Mars.
1278 *Space Science Reviews* 174, 27-48.
- 1279 Michael, P.J., 1988. The concentration, behavior and storage of H₂O in the suboceanic upper
1280 mantle: Implications for mantle metasomatism. *Geochimica et Cosmochimica Acta* 52, 555-566.
- 1281 Mocquet, A., Rosenblatt, P., Dehant, V., Verhoeven, O., 2011. The deep interior of Venus, Mars,
1282 and the Earth: A brief review and the need for planetary surface-based measurements. *Planetary*
1283 *and Space Science* 59, 1048-1061.
- 1284 Mohapatra, R.K., Schwenzer, S.P., Herrmann, S., Murty, S.V.S., Ott, U., Gilmour, J.D., 2009.
1285 Noble gases and nitrogen in Martian meteorites Dar al Gani 476, Sayh al Uhaymir 005 and
1286 Lewis Cliff 88516: EFA and extra neon. *Geochimica et Cosmochimica Acta* 73, 1505-1522.
- 1287 Morschhauser, A., Grott, M., Breuer, D., 2011. Crustal recycling, mantle dehydration, and the
1288 thermal evolution of Mars. *Icarus* 212, 541-558.
- 1289 Musselwhite, D.S., Swindle, T.D., 2001. Is release of martian atmosphere from polar clathrate
1290 the cause of the nakhlite and ALH84001 Ar/Kr/Xe ratios? *Icarus* 154, 207-215.
- 1291 Mysen, B., Boettcher, A., 1975. Melting of a hydrous mantle: I. Phase relations of natural
1292 peridotite at high pressures and temperatures with controlled activities of water, carbon dioxide,
1293 and hydrogen. *Journal of Petrology* 16, 520-548.
- 1294 Mysen, B.O., Cody, G.D., 2004. Solubility and solution mechanism of H₂O in alkali silicate
1295 melts and glasses at high pressure and temperature. *Geochimica et Cosmochimica Acta* 68,
1296 5113-5126.

- 1297 Mysen, B.O., Cody, G.D., Smith, A., 2004. Solubility mechanisms of fluorine in peralkaline and
1298 meta-aluminous silicate glasses and in melts to magmatic temperatures. *Geochimica et*
1299 *Cosmochimica Acta* 68, 2745-2769.
- 1300 Mysen, B.O., Virgo, D., Popp, R.K., Bertka, C.M., 1998. The role of H₂O in Martian magmatic
1301 systems. *American Mineralogist* 83, 942-946.
- 1302 Nekvasil, H., Dondolini, A., Horn, J., Filiberto, J., Long, H., Lindsley, D.H., 2004. The origin
1303 and evolution of silica-saturated alkalic suites: an experimental study. *Journal of Petrology* 45,
1304 693-721.
- 1305 Nekvasil, H., Filiberto, J., McCubbin, F.M., Lindsley, D.H., 2007. Alkalic parental magmas for
1306 the chassignites? *Meteoritics & Planetary Science* 42, 979-992.
- 1307 Nekvasil, H., McCubbin, F.M., Harrington, A.D., Elardo, S.M., Lindsley, D.H., 2009. Linking
1308 the Chassigny meteorite and the Martian surface rock Backstay: Insights into igneous crustal
1309 differentiation processes on Mars. *Meteoritics & Planetary Science* 44, 853-869.
- 1310 Norman, M., 1999. The composition and thickness of the crust of Mars estimated from REE and
1311 Nd isotopic compositions of Martian meteorites. *Meteoritics & Planetary Science* 34, 439-449.
- 1312 Norris, J.R., Herd, C.D.K., 2006. The Yamato 980459 Liquidus at 10 to 20 Kilobars. *Lunar and*
1313 *Planetary Science XXXVII*, Abstract #1787.
- 1314 Nyquist, L.E., Shih, C.Y., McCubbin, F.M., Santos, A.R., Shearer Jr, C.K., Peng, Z.X., Burger,
1315 P.V., Agee, C.B., 2016. Rb-Sr and Sm-Nd isotopic and REE studies of igneous components in
1316 the bulk matrix domain of martian breccia Northwest Africa 7034. . *Meteoritics and Planetary*
1317 *Science* 51, 483-498.
- 1318 O'Reilly, S.Y., Griffin, W.L., 2000. Apatite in the mantle: implications for metasomatic
1319 processes and high heat production in Phanerozoic mantle. *Lithos* 53, 217-232.
- 1320 Ogawa, M., Yanagisawa, T., 2012. Two-dimensional numerical studies on the effects of water on
1321 Martian mantle evolution induced by magmatism and solid-state mantle convection. *Journal of*
1322 *Geophysical Research: Planets* (1991–2012) 117.
- 1323 Ott, U., 1988. Noble gases in SNC meteorites: Shergotty, Nakhla, Chassigny. *Geochimica et*
1324 *Cosmochimica Acta* 52, 1937-1948.
- 1325 Ott, U., 2008. An almost infinite sink for tightly bound xenon: etched Shergotty and (less so)
1326 etched Nakhla. *Lunar and Planetary Science Conference 39th*, Abstract #1096.
- 1327 Owen, T., Biemann, K., Rushneck, D.R., Biller, J.E., Howarth, D.W., Lafleur, A.L., 1977. The
1328 composition of the atmosphere at the surface of Mars. *Journal of Geophysical Research* 82,
1329 6435-6439.
- 1330 Owen, T., Maillard, J.P., De Bergh, C., Lutz, B.L., 1988. Deuterium on Mars: The abundance of
1331 HDO and the value of D/H. *Science* 240, 1767.

- 1332 Ozima, M., Podosek, F.A., 2002. Noble gas geochemistry. Cambridge University Press.
- 1333 Papike, J.J., Burger, P.V., Shearer, C.K., McCubbin, F.M., 2013. Experimental and crystal
1334 chemical study of the basalt–eclogite transition in Mars and implications for martian magmatism.
1335 *Geochimica et Cosmochimica Acta* 104, 358-376.
- 1336 Parman, S.W., Grove, T.L., Kelley, K.A., Plank, T., 2011. Along-Arc Variations in the Pre-
1337 Eruptive H₂O Contents of Mariana Arc Magmas Inferred from Fractionation Paths. *Journal of*
1338 *Petrology* 52, 257-278.
- 1339 Patiño Douce, A.E., Roden, M., 2006. Apatite as a probe of halogen and water fugacities in the
1340 terrestrial planets. *Geochimica et Cosmochimica Acta* 70, 3173-3196.
- 1341 Patiño Douce, A.E., Roden, M.F., Chaumba, J., Fleisher, C., Yogodzinski, G., 2011.
1342 Compositional variability of terrestrial mantle apatites, thermodynamic modeling of apatite
1343 volatile contents, and the halogen and water budgets of planetary mantles. *Chemical Geology*
1344 288, 14-31.
- 1345 Pauer, M., Breuer, D., 2008. Constraints on the maximum crustal density from gravity–
1346 topography modeling: Applications to the southern highlands of Mars. *Earth and Planetary*
1347 *Science Letters* 276, 253-261.
- 1348 Pawley, A.R., Holloway, J.R., McMillan, P.F., 1992. The effect of oxygen fugacity on the
1349 solubility of carbon-oxygen fluids in basaltic melt. *Earth and Planetary Science Letters* 110, 213-
1350 225.
- 1351 Pepin, R.O., 1991. On the origin and early evolution of terrestrial planet atmospheres and
1352 meteoritic volatiles. *Icarus* 92, 2-79.
- 1353 Perron, J.T., Mitrovica, J.X., Manga, M., Matsuyama, I., Richards, M.A., 2007. Evidence for an
1354 ancient martian ocean in the topography of deformed shorelines. *Nature* 447, 840-843.
- 1355 Peters, T.J., Simon, J.I., Jones, J.H., Usui, T., Moriwaki, R., Economos, R.C., Schmitt, A.K.,
1356 McKeegan, K.D., 2015. Tracking the source of the enriched martian meteorites in olivine-hosted
1357 melt inclusions of two depleted shergottites, Yamato 980459 and Tissint. *Earth and Planetary*
1358 *Science Letters* 418, 91-102.
- 1359 Piccoli, P.M., Candela, P.A., 2002. Apatite in igneous systems. *Reviews in Mineralogy and*
1360 *Geochemistry* 48, 255-292.
- 1361 Plesa, A.-C., Breuer, D., 2014. Partial melting in one-plate planets: implications for thermo-
1362 chemical and atmospheric evolution. *Planetary and Space Science* 98, 50-65.
- 1363 Plesa, A.C., Tosi, N., Breuer, D., 2014. Can a fractionally crystallized magma ocean explain the
1364 thermo-chemical evolution of Mars? *Earth and Planetary Science Letters* 403, 225-235.
- 1365 Pommier, A., Grove, T.L., Charlier, B., 2012. Water storage and early hydrous melting of the
1366 Martian mantle. *Earth and Planetary Science Letters* 333–334, 272-281.

1367 Poulet, F., Bibring, J.P., Mustard, J.F., Gendrin, A., Mangold, N., Langevin, Y., Arvidson, R.E.,
 1368 Gondet, B., Gomez, C., 2005. Phyllosilicates on Mars and implications for early martian climate.
 1369 Nature 438, 623-627.

1370 Richter, K., Pando, K., Danielson, L.R., 2009. Experimental evidence for sulfur-rich martian
 1371 magmas: Implications for volcanism and surficial sulfur sources. Earth and Planetary Science
 1372 Letters 288, 235-243.

1373 Richter, K., Yang, H., Costin, G., Downs, R.T., 2008. Oxygen fugacity in the Martian mantle
 1374 controlled by carbon: New constraints from the nakhlite MIL 03346. Meteoritics & Planetary
 1375 Science 43, 1709-1723.

1376 Roberts, J.H., Lillis, R.J., Manga, M., 2009. Giant impacts on early Mars and the cessation of the
 1377 Martian dynamo. Journal of Geophysical Research: Planets 114, n/a-n/a.

1378 Rogers, A.D., Nekvasil, H., 2015. Feldspathic rocks on Mars: Compositional constraints from
 1379 infrared spectroscopy and possible formation mechanisms. Geophysical Research Letters 42,
 1380 2619-2626.

1381 Rossi, M., Ghiara, M.R., Chita, G., Capitelli, F., 2011. Crystal-chemical and structural
 1382 characterization of fluorapatites in ejecta from Somma-Vesuvius volcanic complex. American
 1383 Mineralogist 96, 1828-1837.

1384 Ruedas, T., Tackley, P.J., Solomon, S.C., 2013. Thermal and compositional evolution of the
 1385 martian mantle: Effects of water. Physics of The Earth and Planetary Interiors 220, 50-72.

1386 Saal, A.E., Hauri, E.H., Langmuir, C.H., Perfit, M.R., 2002. Vapour undersaturation in primitive
 1387 mid-ocean-ridge basalt and the volatile content of Earth's upper mantle. Nature 419, 451-455.

1388 Salters, V.J.M., Longhi, J.E., Bizimis, M., 2002. Near mantle solidus trace element partitioning
 1389 at pressures up to 3.4 GPa. Geochemistry, Geophysics, Geosystems 3, 1-23.

1390 Sandland, T.O., Du, L.-S., Stebbins, J.F., Webster, J.D., 2004. Structure of Cl-containing silicate
 1391 and aluminosilicate glasses: A ³⁵Cl MAS-NMR study. Geochimica et Cosmochimica Acta 68,
 1392 5059-5069.

1393 Sandu, C., Kiefer, W.S., 2012. Degassing history of Mars and the lifespan of its magnetic
 1394 dynamo. Geophysical Research Letters 39, 10.1029/2011GL050225.

1395 Santos, A.R., Agee, C.B., McCubbin, F.M., Shearer, C.K., Burger, P.V., Tartèse, R., Anand, M.,
 1396 2015. Petrology of igneous clasts in Northwest Africa 7034: Implications for the petrologic
 1397 diversity of the martian crust. Geochimica et Cosmochimica Acta 157, 56-85.

1398 Sautter, V., Fabre, C., Forni, O., Toplis, M.J., Cousin, A., Ollila, A.M., Meslin, P.Y., Maurice,
 1399 S., Wiens, R.C., Baratoux, D., Mangold, N., Le Mouélic, S., Gasnault, O., Berger, G., Lasue, J.,
 1400 Anderson, R.A., Lewin, E., Schmidt, M., Dyar, D., Ehlmann, B.L., Bridges, J., Clark, B., Pinet,
 1401 P., 2014. Igneous mineralogy at Bradbury Rise: The first ChemCam campaign at Gale crater.
 1402 Journal of Geophysical Research: Planets 119, 2013JE004472.

- 1403 Sautter, V., Jambon, A., Boudouma, O., 2006. Cl-amphibole in the nakhlite MIL 03346:
1404 Evidence for sediment contamination in a Martian meteorite. *Earth and Planetary Science Letters*
1405 252, 45-55.
- 1406 Sautter, V., Toplis, M.J., Wiens, R.C., Cousin, A., Fabre, C., Gasnault, O., Maurice, S., Forni,
1407 O., Lasue, J., Ollila, A., Bridges, J.C., Mangold, N., Le Mouelic, S., Fisk, M., Meslin, P.Y.,
1408 Beck, P., Pinet, P., Le Deit, L., Rapin, W., Stolper, E.M., Newsom, H., Dyar, D., Lanza, N.,
1409 Vaniman, D., Clegg, S., Wray, J.J., 2015. In situ evidence for continental crust on early Mars.
1410 *Nature Geosci* 8, 605-609.
- 1411 Schaller, T., Dingwell, D.B., Keppler, H., Knöller, W., Merwin, L., Sebal, A., 1992. Fluorine in
1412 silicate glasses: A multinuclear nuclear magnetic resonance study. *Geochimica et Cosmochimica*
1413 *Acta* 56, 701-707.
- 1414 Scheinberg, A., Elkins-Tanton, L.T., Zhong, S.J., 2014. Timescale and morphology of Martian
1415 mantle overturn immediately following magma ocean solidification. *Journal of Geophysical*
1416 *Research: Planets* 119, 454-467.
- 1417 Schwenzer, S., Greenwood, R., Kelley, S., Ott, U., Tindle, A., Haubold, R., Herrmann, S.,
1418 Gibson, J., Anand, M., Hammond, S., 2013. Quantifying noble gas contamination during
1419 terrestrial alteration in Martian meteorites from Antarctica. *Meteoritics & Planetary Science* 48,
1420 929-954.
- 1421 Schwenzer, S., Ott, U., 2006. Evaluating Kr-and Xe-Data in the Nakhilites and ALHA84001-
1422 Does EFA Hide EFM?, 37th Annual Lunar and Planetary Science Conference, p. 1614.
- 1423 Schwenzer, S.P., Bullock, M.A., Bridges, J., Chavez, C., Filiberto, J., Hicks, L.J., Kelley, S.P.,
1424 Miller, M.A., Moore, J.M., Smith, H., Swindle, T.D., treiman, A.H., 2016. Noble Gas
1425 Fractionation in Hydrous Rock Alteration under Diagenetic Pressure and Temperature
1426 Conditions. 47th Lunar and Planetary Science Conference, Abstract #1889.
- 1427 Schwenzer, S.P., Herrmann, S., Mohapatra, R.K., Ott, U., 2007. Noble gases in mineral separates
1428 from three shergottites: Shergotty, Zagami, and EETA79001. *Meteoritics & Planetary Science*
1429 42, 387-412.
- 1430 Shahr, A., Hillgren, V., Horan, M., Mesa-Garcia, J., Kaufman, L., Mock, T., 2015. Sulfur-
1431 controlled iron isotope fractionation experiments of core formation in planetary bodies.
1432 *Geochimica et Cosmochimica Acta* 150, 253-264.
- 1433 Sharp, Z.D., McCubbin, F.M., Shearer, C.K., 2013. A hydrogen-based oxidation mechanism
1434 relevant to planetary formation. *Earth and Planetary Science Letters* 380, 88-97.
- 1435 Sharp, Z.D., Williams, J., Shearer, C.K., Agee, C.B., McKeegan, K.D., 2016. The Chlorine
1436 Isotope Composition of Martian Meteorites 2. Implications for the early Solar System and the
1437 formation of Mars. *Meteoritics and Planetary Science* this volume.

- 1438 Shearer, C., Burger, P., Papike, J., McCubbin, F., Bell, A., 2015. Crystal chemistry of merrillite
1439 from Martian meteorites: Mineralogical recorders of magmatic processes and planetary
1440 differentiation. *Meteoritics & Planetary Science* 50, 649-673.
- 1441 Silver, L.A., Ihinger, P.D., Stolper, E., 1990. The influence of bulk composition on the speciation
1442 of water in silicate glasses. *Contributions to Mineralogy and Petrology* 104, 142-162.
- 1443 Spengler, S.R., Garcia, M.O., 1988. Geochemistry of the Hawi Lavas, Kohala Volcano, Hawaii.
1444 *Contributions to Mineralogy and Petrology* 99, 90-104.
- 1445 Stanley, B.D., Hirschmann, M.M., Withers, A.C., 2011. CO₂ solubility in Martian basalts and
1446 Martian atmospheric evolution. *Geochimica et Cosmochimica Acta* 75, 5987-6003.
- 1447 Stanley, B.D., Hirschmann, M.M., Withers, A.C., 2014. Solubility of COH volatiles in graphite-
1448 saturated martian basalts. *Geochimica et Cosmochimica Acta* 129, 54-76.
- 1449 Stanley, B.D., Schaub, D.R., Hirschmann, M.M., 2012. CO₂ solubility in primitive martian
1450 basalts similar to Yamato 980459, the effect of composition on CO₂ solubility of basalts, and the
1451 evolution of the martian atmosphere. *American Mineralogist* 97, 1841-1848.
- 1452 Steele, A., Fries, M.D., Amundsen, H.E.F., Mysen, B.O., Fogel, M.L., Schweizer, M., Bockor,
1453 N.Z., 2007. Comprehensive imaging and Raman spectroscopy of carbonate globules from
1454 Martian meteorite ALH 84001 and a terrestrial analogue from Svalbard. *Meteoritics & Planetary
1455 Science* 42, 1549-1566.
- 1456 Steele, A., McCubbin, F.M., Fries, M., Kater, L., Bockor, N.Z., Fogel, M.L., Conrad, P.G.,
1457 Glamoclija, M., Spencer, M., Morrow, A.L., Hammond, M.R., Zare, R.N., Vicenzi, E.P.,
1458 Siljeström, S., Bowden, R., Herd, C.D.K., Mysen, B.O., Shirey, S.B., Amundsen, H.E.F.,
1459 Treiman, A.H., Bullock, E.S., Jull, A.J.T., 2012a. A Reduced Organic Carbon Component in
1460 Martian Basalts. *Science* 337, 212-215.
- 1461 Steele, A., McCubbin, F.M., Fries, M.D., 2016. The provenance and formation of reduced
1462 carbon phases in martian meteorites. *Meteoritics and Planetary Science* this volume.
- 1463 Steele, A., McCubbin, F.M., Fries, M.D., Golden, D.C., Ming, D.W., Benning, L.G., 2012b.
1464 Graphite in the martian meteorite Allan Hills 84001. *American Mineralogist* 97, 1256-1259.
- 1465 Stewart, A.J., Schmidt, M.W., van Westrenen, W., Liebske, C., 2007. Mars: A New Core-
1466 Crystallization Regime. *Science* 316, 1323-1325.
- 1467 Stoppa, F., Liu, Y., 1995. Chemical composition and petrogenetic implications of apatites from
1468 some ultra-alkaline Italian rocks. *European Journal of Mineralogy* 7, 391-402.
- 1469 Swindle, T., Grier, J., Burkland, M., 1995. Noble gases in orthopyroxenite ALH84001: A
1470 different kind of martian meteorite with an atmospheric signature. *Geochimica et Cosmochimica
1471 Acta* 59, 793-801.

- 1472 Swindle, T., Thomas, C., Mousis, O., Lunine, J., Picaud, S., 2009. The trapping of Ar, Kr, and
1473 Xe in martian clathrates and the possibility of detecting clathrates on Mars by seasonal changes
1474 in the Xe/Kr ratio. Lunar and Planetary Science Conference 40, abstract# 1660.
- 1475 Swindle, T.D., 2002. Martian Noble Gases. Reviews in Mineralogy and Geochemistry 47, 171-
1476 190.
- 1477 Taylor, G.J., 2013. The bulk composition of Mars. Chemie der Erde - Geochemistry, 401-420.
- 1478 Taylor, G.J., Boynton, W.V., Brückner, J., Wänke, H., Dreibus, G., Kerry, K., Keller, J., Reedy,
1479 R., Evans, L., Starr, R., Squyres, S., Karunatillake, S., Gasnault, O., Maurice, S., d'Uston, C.,
1480 Englert, P., Dohm, J., Baker, V., Hamara, D., Janes, D., Sprague, A., Kim, K., Drake, D., 2006.
1481 Bulk composition and early differentiation of Mars. Journal of Geophysical Research 111,
1482 doi:10.1029/2005JE002645.
- 1483 Taylor, S.R., 2001. Solar System Evolution: A new perspective 2nd edition, 2nd Edition ed.
1484 Cambridge University Press, Canberra, Australia.
- 1485 Tian, F., Kasting, J.F., Solomon, S.C., 2009. Thermal escape of carbon from the early Martian
1486 atmosphere. Geophysical Research Letters 36.
- 1487 Treiman, A.H., 1985. Amphibole and hercynite spinel in Shergotty and Zagami Magmatic water,
1488 depth of crystallization, and metasomatism. Meteoritics 20, 229-243.
- 1489 Treiman, A.H., Dyar, M.D., McCanta, M., Noble, S.K., Pieters, C.M., 2007. Martian Dunit
1490 NWA 2737: Petrographic constraints on geological history, shock events, and olivine color.
1491 Journal of Geophysical Research 112, doi:10.1029/2006JE002777.
- 1492 Usui, T., Alexander, C.M.O.D., Wang, J., Simon, J.I., Jones, J.H., 2012. Origin of water and
1493 mantle–crust interactions on Mars inferred from hydrogen isotopes and volatile element
1494 abundances of olivine-hosted melt inclusions of primitive shergottites. Earth and Planetary
1495 Science Letters 357–358, 119-129.
- 1496 Usui, T., Alexander, C.M.O.D., Wang, J., Simon, J.I., Jones, J.H., 2015. Meteoritic evidence for
1497 a previously unrecognized hydrogen reservoir on Mars. Earth and Planetary Science Letters 410,
1498 140-151.
- 1499 Villanueva, G.L., Mumma, M.J., Novak, R.E., Käufl, H.U., Hartogh, P., Encrenaz, T., Tokunaga,
1500 A., Khayat, A., Smith, M.D., 2015. Strong water isotopic anomalies in the martian atmosphere:
1501 Probing current and ancient reservoirs. Science 348, 218-221.
- 1502 Wallace, P.J., Edmonds, M., 2011. The Sulfur Budget in Magmas: Evidence from Melt
1503 Inclusions, Submarine Glasses, and Volcanic Gas Emissions. Reviews in Mineralogy and
1504 Geochemistry 73, 215-246.
- 1505 Wänke, H., 1981. Constitution of Terrestrial Planets. Philosophical Transactions of the Royal
1506 Society of London Series a-Mathematical Physical and Engineering Sciences 303, 287-302.

- 1507 Wänke, H., 1991. Chemistry, Accretion, and Evolution of Mars. *Space Science Reviews* 56, 1-8.
- 1508 Wänke, H., Dreibus, G., 1988. Chemical-Composition and Accretion History of Terrestrial
1509 Planets. *Philosophical Transactions of the Royal Society of London Series a-Mathematical*
1510 *Physical and Engineering Sciences* 325, 545-557.
- 1511 Watson, L.L., Epstein, S., Stolper, E.M., 1994a. D/H of Water Released by Stepped Heating of
1512 Shergotty, Zagami, Chassigny, Alh-84001, and Nakhla. *Meteoritics* 29, 547-547.
- 1513 Watson, L.L., Hutcheon, I.D., Epstein, S., Stolper, E.M., 1994b. Water on Mars: Clues from
1514 Deuterium/Hydrogen and Water Contents of Hydrous Phases in SNC Meteorites. *Science* 265,
1515 86-90.
- 1516 Webster, C.R., Mahaffy, P.R., Flesch, G.J., Niles, P.B., Jones, J.H., Leshin, L.A., Atreya, S.K.,
1517 Stern, J.C., Christensen, L.E., Owen, T., 2013. Isotope ratios of H, C, and O in CO₂ and H₂O of
1518 the martian atmosphere. *Science* 341, 260-263.
- 1519 Webster, J., 1997. Chloride Solubility in Felsic Melts and the Role of Chloride in Magmatic
1520 Degassing. *Journal of Petrology* 38, 1793-1807.
- 1521 Webster, J.D., 1990. Partitioning of F between H₂O and CO₂ fluids and topaz rhyolite melt.
1522 *Contributions to Mineralogy and Petrology* 104, 424-438.
- 1523 Webster, J.D., De Vivo, B., 2002. Experimental and modeled solubilities of chlorine in
1524 aluminosilicate melts, consequences of magma evolution, and implications for exsolution of
1525 hydrous chloride melt at Mt. Somma-Vesuvius. *American Mineralogist* 87, 1046-1061.
- 1526 Webster, J.D., Kinzler, R.J., Mathez, E.A., 1999. Chloride and water solubility in basalt and
1527 andesite melts and implications for magmatic degassing. *Geochimica et Cosmochimica Acta* 63,
1528 729-738.
- 1529 Webster, J.D., Vetere, F., Botcharnikov, R.E., Goldoff, B., McBirney, A., Doherty, A.L., 2015.
1530 Experimental and modeled chlorine solubilities in aluminosilicate melts at 1 to 7000 bars and
1531 700 to 1250 °C: Applications to magmas of Augustine Volcano, Alaska. *American Mineralogist*
1532 100, 522-535.
- 1533 Whitaker, M., Nekvasil, H., Lindsley, D., McCurry, M., 2008. Can crystallization of olivine
1534 tholeiite give rise to potassic rhyolites?—an experimental investigation. *Bulletin of Volcanology*
1535 70, 417-434.
- 1536 Wiens, R.C., 1988. Noble gases released by vacuum crushing of EETA 79001 glass. *Earth and*
1537 *Planetary Science Letters* 91, 55-65.
- 1538 Williams, J.T., Shearer Jr, C.K., Sharp, Z.D., Burger, P.V., McCubbin, F.M., Santos, A.R., Agee,
1539 C.B., McKeegan, K.D., 2016. The chlorine isotopic composition of martian meteorites 1.
1540 Chlorine isotopic composition of the martian mantle, crustal, and atmospheric reservoirs and
1541 their interactions. *Meteoritics & Planetary Science*. this volume.

- 1542 Wray, J.J., Hansen, S.T., Dufek, J., Swayze, G.A., Murchie, S.L., Seelos, F.P., Skok, J.R., Irwin
1543 Iii, R.P., Ghiorso, M.S., 2013. Prolonged magmatic activity on Mars inferred from the detection
1544 of felsic rocks. *Nature Geosci* 6, 1013-1017.
- 1545 Zeng, Q., Nekvasil, H., Grey, C.P., 1999. Proton environments in hydrous aluminosilicate
1546 glasses: A H-1 MAS, H-1/Al-27, and H-1/Na-23 TRAPDOR NMR study. *Journal of Physical*
1547 *Chemistry B* 103, 7406-7415.
- 1548 Zeng, Q., Stebbins, J.F., 2000. Fluoride sites in aluminosilicate glasses; high-resolution 19 F
1549 NMR results. *American Mineralogist* 85, 863-867.
- 1550 Zimova, M., Webb, S., 2006. The effect of chlorine on the viscosity of Na₂O-Fe₂O₃-Al₂O₃-SiO₂
1551 melts. *American Mineralogist* 91, 344-352.
- 1552 Zimova, M., Webb, S.L., 2007. The combined effects of chlorine and fluorine on the viscosity of
1553 aluminosilicate melts. *Geochimica et Cosmochimica Acta* 71, 1553-1562.
1554
1555




Figure 1. Ternary plot of apatite X-site occupancy (mol%) from shergottites (red) and terrestrial basalts (blue). The shergottite data include basaltic, lherzolithic, and olivine-phyric shergottites, and data are from Gross et al. (2013) and McCubbin et al. (submitted; 2012). Terrestrial apatites are from basaltic systems that have not been reported to have interacted with hydrothermal fluids (Brown and Peckett 1977; Spengler and Garcia 1988; Liu and Comodi 1993; Stoppa and Liu 1995; Coulson and Chambers 1996; O'Reilly and Griffin 2000; Dymek and Owens 2001; Piccoli and Candela 2002; Casey et al. 2007; Filiberto and Treiman 2009b; Rossi et al. 2011).




Figure 2: Hydrogen isotopic compositions of Martian water reservoirs, modified and updated from Usui et al. (2012). Data sources: Martian mantle (Chassigny: Leshin et al., 1996, Watson et al., 1994) Depleted Shergottites; Y980459 from Usui et al., 2012; Tissint from Mane et al., submitted. Nakhrites: Nakhla from Hallis et al., (2012); NWA 817 from Gillet et al., (2002). Intermediate reservoir: EETA 79001 (A) from Usui et al., (2015); Clays from Yellowknife Bay from Mahaffy et al., (2015). Atmospheric reservoir (Leshin et al., 2013; Webster et al., 2013).




Figure 3. Schematic diagram locating the different gas reservoirs found in Martian meteorites and processes shaping them. For endmembers Martian atmosphere from Bogard and Johnson (1983), Chassigny from Ott (1988), but see also Swindle (2002). Terrestrial atmosphere from (Ozima and Podosek 2002). Earth atmosphere is fractionated when dissolved in water (e.g. Ozima and Podosek 2002 for discussion) but also when incorporated in minerals (Mohapatra et al. 2009; Schwenger et al. 2013). The meteorites show the following signatures: Chassigny noble gases are thought to represent the Martian interior (e.g., Ott 1988, Matthew and Marti 2001), the Shergottites a mixture of implanted Martian atmosphere and Martian interior (often with some admixed terrestrial air contamination; Bogard and Johnson 1983, Ott 1988, Swindle 2002), and the nakhlites and ALH84001 (Ott 1988, Swindle 2002, and see text). NWA 7034 from

1584 Cartwright et al. (2014). Meteorite measurements often return a mixture of two or more
1585 components as indicated by the red arrows. The ‘crush’ component was found by Wiens (1988)
1586 and has not been understood yet.

1587






Figure 4. Liquidus depression ($\Delta T^{\circ}\text{C}$) as a function of wt% water in the melt from terrestrial experiments over a range of pressures and experimental techniques Medard and Grove (2008); Almeev et al. (2007); Katz et al. (2003); and Danyushevsky (2001). Also, plotted is the calculated liquidus depression (red curve) for experiments on the Martian meteorite Yamato 980459 with <0.5, 0.5, and 2 wt% water (Norris and Herd 2006; Dalton et al. 2007; Draper 2007 respectively) calculated relative to the anhydrous experiments of Musselwhite et al. (2006).





1596


1597 **Figure 5.** An example calculation of mantle degassing and water loss on Mars, shown as a
1598 function of time since accretion. In this simulation, two-thirds of the original mantle water is lost
1599 from the mantle, primarily in the first billion years of Martian history.

1600









Mantle Water Loss

